PRIOR AUTHORIZATION CRITERIA

This list is current as of 10/1/2025 and pertains to the following formularies:

2026 Independent Health's Medicare Advantage Enhanced Part D Formulary 2026 Independent Health's Medicare Advantage Employer Group's Part D Formulary

Independent Health requires you (or your physician) to get prior authorization for certain drugs. This means that you will need to get approval from us before you fill your prescriptions. If you do not get approval, we may not cover the drug. These drugs are listed with a "PA" in the Requirements/Limits column on the formularies. This document contains the Prior Authorization requirements that are associated with our Medicare Advantage Part D formularies.

If you have any questions, please contact our Medicare Member Services Department at 1-800-665-1502 or, for TTY users 711, October 1st – March 31st: Monday through Sunday from 8 a.m. to 8 p.m., April 1st – September 30th: Monday through Friday from 8 a.m. to 8 p.m.

The formulary may change at any time. You will receive notice when necessary.

ACTIMMUNE (interferon gamma-1b)

Products Affected

• ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ADEMPAS (riociguat)

Products Affected

 ADEMPAS ORAL TABLET 0.5 MG, 1 MG, 1.5 MG, 2 MG, 2.5 MG

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, severe hepatic impairment (Child-Pugh class C), creatinine clearance below 15 mL/min or on dialysis, concurrent use with nitrates or nitric oxide donors in any form, concurrent use with phosphodiesterase inhibitors
Required Medical Information	Diagnosis of covered use confirmed by right heart catheterization, submission of patient's WHO Group classification, mean pulmonary arterial pressure greater than 20 mm Hg at rest, pulmonary arterial wedge pressure less than or equal to 15 mm Hg, creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), and pregnancy status for female patients of childbearing potential. For pulmonary arterial hypertension (PAH, WHO Group 1), documentation of pulmonary vascular resistance (PVR) greater than 2 Woods units, submission of current or previous therapies used to treat the condition (see Other Criteria). For chronic thromboembolic pulmonary hypertension (CTEPH, WHO Group 4), confirmation of PVR greater than 3 Woods units, evidence of chronic pulmonary embolism on computed tomography or ventilation/perfusion (V/Q) scan, and attestation patient has inoperable disease or has persistent or recurrent disease after CTEPH surgery (pulmonary thromboendarterectomy).
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology and pulmonology
Coverage Duration	1 year
Other Criteria	For initial authorization for PAH (WHO Group 1), the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to both (1) sildenafil or tadalafil and (2) ambrisentan or bosentan.

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

ALECENSA (alectinib)

Products Affected

• ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ALUNBRIG (brigatinib)

Products Affected

• ALUNBRIG

PA Criteria	Criteria Details
Exclusion Criteria	Uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, baseline blood pressure reading, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

AUGTYRO (repotrectinib)

Products Affected

• AUGTYRO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inhibitors or inducers or P-glycoprotein inhibitors
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For non-small cell lung cancer, submission of test confirming tumor is ROS1-positive. For other solid tumors, submission of test confirming tumor has a neurotrophic tyrosine receptor kinase (NTRK) gene fusion, attestation patient has progressed following treatment or patient has no satisfactory alternative therapy.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

AUVELITY (dextromethorphan/bupropion)

Products Affected

• AUVELITY

PA Criteria	Criteria Details
Exclusion Criteria	Seizure disorder, current or prior diagnosis of bulimia or anorexia nervosa, severe hepatic impairment, severe renal impairment, administration of monoamine oxidase inhibitors within 14 days of initiation
Required Medical Information	Diagnosis of covered use, attestation patient has been screened for and does not have bipolar disorder, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to psychiatry
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to two generic on-formulary antidepressants (e.g., bupropion, SSRI, SNRI).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

AYVAKIT (avapritinib)

Products Affected

 AYVAKIT ORAL TABLET 100 MG, 200 MG, 25 MG, 300 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers or strong CYP3A inhibitors. For advanced or indolent systemic mastocytosis, platelet count below 50 x 10^9/L.
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For gastrointestinal stromal tumor (GIST), submission of test result confirming presence of PDGFRA exon 18 mutation. For advanced or indolent systemic mastocytosis, submission of platelet count.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to allergy, hematology, immunology, and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

BALVERSA (erdafitinib)

Products Affected

• BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	PD-1/PD-L1 inhibitor-eligible patients who have not received this therapy, coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of susceptible FGFR3 genetic alterations, submission of current or previous therapies used to treat the condition (see Other Criteria), pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	This drug is not recommended for the treatment of patients who are eligible for and have not received prior PD-1 or PD-L1 inhibitor therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

BEMPEDOIC ACID

Products Affected

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant pravastatin utilization with doses above 40 mg/day, concomitant simvastatin utilization with doses above 20 mg/day, history of tendon disorders or rupture
Required Medical Information	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must (1) currently be using a statin (unless contraindicated) plus ezetimibe, or (2) have tried and failed to have an adequate response to or had an intolerance to (a) at least two statins or (b) one statin and ezetimibe. At least one statin previously tried and failed must be a hydrophilic statin. For each annual reauthorization, documentation that the patient remains on previously-used lipid-lowering therapies since the previous approval, unless there is documentation of a new contraindication or intolerance requiring discontinuation of a therapy (or therapies) since the previous approval, is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

BENLYSTA (belimumab)

Products Affected

• BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	Severe active central nervous system lupus, patients using other biologic medications or intravenous cyclophosphamide
Required Medical Information	Diagnosis of covered use, submission of current therapies used to treat the condition (see Other Criteria). For systemic lupus erythematosus, submission of autoantibody-positive test result for anti-nuclear antibodies (ANA) and/or anti-double-stranded DNA (anti-dsDNA).
Age Restrictions	
Prescriber Restrictions	Restricted to immunology, nephrology, and rheumatology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must be using standard therapy, defined as at least one of the following: systemic corticosteroids (e.g., prednisone), antimalarials (e.g., hydroxychloroquine), or immunosuppressants (e.g., azathioprine, methotrexate, mycophenolate mofetil). For each annual reauthorization, confirmation patient is still using some form of standard therapy (as defined above), unless contraindicated, is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

BESREMI (ropeginterferon alfa-2b-njft)

Products Affected

• BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	History or presence of severe psychiatric disorders (including severe depression or suicidal ideation), history of presence of active serious or untreated autoimmune disease, moderate or severe hepatic impairment (Child-Pugh class B or C), immunosuppressed transplant recipients, severe or unstable cardiovascular disease (e.g., uncontrolled hypertension, NYHA class 2-4 congestive heart failure, serious cardiac arrhythmia, significant coronary artery stenosis, unstable angina), stroke or myocardial infarction within previous 6 months, severe renal impairment (eGFR less than 30 mL/min)
Required Medical Information	Diagnosis of covered use, submission of eGFR, documentation patient has tried and failed to have an adequate response to or had an intolerance/contraindication to hydroxyurea (HU), pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	If patient is taking HU, initially 12 weeks, then 1 year. If patient is not taking HU, 1 year.
Other Criteria	For the first reauthorization in patients using HU at the start of therapy, attestation patient has tapered completely off HU by the end of week 12 is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

BEXAROTENE GEL

Products Affected

• bexarotene external

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

BIOLOGIC RESPONSE MODIFIERS

Products Affected

- KINERET SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- OTEZLA ORAL TABLET 30 MG
- OTEZLA ORAL TABLET THERAPY PACK 10 & 20 & 30 MG
- SOTYKTU
- STELARA SUBCUTANEOUS SOLUTION 45

MG/0.5ML

- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- tyenne subcutaneous
- ustekinumab subcutaneous
- VELSIPITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria). For all drugs managed by this policy except Otezla and Velsipity, submission of baseline latent tuberculosis screening test (Mantoux tuberculin skin test [a.k.a. PPD test] or interferon-gamma release assay [IGRA]).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For initial authorization of a drug managed by this policy, the patient must have tried and failed to have an adequate response to or had an intolerance to at least two preferred agents (an adalimumab biosimilar, Cosentyx, Enbrel, Rinvoq, Skyrizi, an ustekinumab biosimilar, and Xeljanz/Xeljanz XR) for the indication submitted, where possible. For all drugs managed by this policy except Otezla and Velsipity, if TB screening test returns a positive result, coverage will be delayed until latent TB is treated. For each annual reauthorization, yearly TB screening test or chest X-ray required for patients who live in, work in, or travel to areas where TB exposure is likely while on treatment or for those who have previously had a positive TB screening test.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

BOSULIF (bosutinib)

Products Affected

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inhibitors or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For accelerated or blast phase Ph+CML, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	For initial authorization for accelerated or blast phase Ph+ CML, the patient must have either (1) tried and had an intolerance to dasatinib, imatinib, or nilotinib or (2) resistance to imatinib, defined as (a) failure to achieve or maintain any hematologic improvement within 4 weeks while on imatinib, or (b) failure to achieve a complete hematologic response by 3 months, cytogenetic response by 6 months or major cytogenetic response by 12 months, or (c) progression of disease after a previous cytogenetic or hematologic response, or (d) presence of a genetic mutation in the BCR-ABL gene associated with imatinib resistance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

BRAFTOVI/MEKTOVI (encorafenib/binimetinib)

Products Affected

- BRAFTOVI ORAL CAPSULE 75 MG
- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of BRAF V600E (or V600K mutation for a melanoma diagnosis, pregnancy status for female patients of childbearing potential. For metastatic melanoma or metastatic non-small cell lung cancer, confirmation that encorafenib and binimetinib will be co-administered. For metastatic colorectal cancer, confirmation that encorafenib and cetuximab will be co-administered.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

BUTALBITAL-CONTAINING PRODUCTS IN OLDER PATIENTS

Products Affected

- ASCOMP-CODEINE
- butalbital-acetaminophen oral tablet 50-325 mg
- butalbital-apap-caff-cod
- butalbital-apap-caffeine oral capsule
- butalbital-apap-caffeine oral tablet 50-325-40 mg
- butalbital-asa-caff-codeine
- butalbital-aspirin-caffeine oral capsule
- TENCON ORAL TABLET 50-325 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation patient has tried and failed a preferred alternative such as ibuprofen or rizatriptan, or has contraindications to all alternatives.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

CABLIVI (caplacizumab-yhdp)

Products Affected

• CABLIVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, confirmation drug will be given with plasma exchange and immunosuppressive therapy. If the coverage determination request is not for the patient's first use of caplacizumab, submission of previous aTTP recurrences while on caplacizumab.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology, hematology, and immunology
Coverage Duration	3 months
Other Criteria	If the coverage determination request is not for the patient's first use of caplacizumab, coverage will not be authorized if the patient has had more than 2 recurrences of aTTP while on therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CALQUENCE (acalabrutinib)

Products Affected

• CALQUENCE ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment, coadministration with strong CYP3A inhibitors
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CAMZYOS (mavacamten)

Products Affected

CAMZYOS

PA Criteria	Criteria Details
Exclusion Criteria	Left ventricular ejection fraction (LVEF) less than 55%, coadministration with strong CYP2C19 inhibitors, moderate or strong CYP2C19 inducers, moderate or strong CYP3A4 inducers, a non-dihydropyridine (DHP) calcium channel blocker (CCB) plus a beta-blocker, disopyramide, or ranolazine
Required Medical Information	Diagnosis of covered use including all three of the following: (1) attestation patient has exertional symptoms consistent with the definition of NYHA class II or III disease, (2) confirmation of left ventricular (LV) outflow tract obstruction gradient of at least 50 mm Hg either at rest, during Valsalva maneuver testing, or after exercise, and (3) confirmation of LV wall thickness of at least 15 mm or at least 13 mm if condition is familial, submission of current LVEF, current or previous therapies used to treat the condition (see Other Criteria), pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to both a beta-blocker and a non-DHP CCB. For each reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

CAPRELSA (vandetanib)

Products Affected

CAPRELSA

PA Criteria	Criteria Details
Exclusion Criteria	History of congenital long QT syndrome, torsades de pointes, uncompensated heart failure, or bradyarrhythmias, QTcF interval greater than 450 msec, hypocalcemia, hypokalemia, hypomagnesemia, coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of baseline serum potassium, calcium, magnesium, creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), ECG (or QT/QTcF interval), and pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CARGLUMIC ACID

Products Affected

• carglumic acid oral tablet soluble

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of elevated plasma ammonia level.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated plasma ammonia level since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CERDELGA (eliglustat)

Products Affected

• CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	Ultrarapid CYP2D6 metabolizers, pre-existing cardiac disease, moderate or severe hepatic impairment, long QT syndrome, coadministration with Class Ia or Class III antiarrhythmics. In patients who are poor or intermediate CYP2D6 metabolizers only, mild hepatic impairment.
Required Medical Information	Diagnosis of covered use, submission of CYP2D6 metabolizer status as detected by a test for determining CYP2D6 genotype, liver function testing or Child-Pugh score.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated liver function testing or Child- Pugh score since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CFTR MODULATOR THERAPIES

Products Affected

- KALYDECO
- ORKAMBI
- SYMDEKO

50-75 & 150 MG, 50-25-37.5 & 75 MG

• TRIKAFTA ORAL THERAPY PACK

• TRIKAFTA ORAL TABLET THERAPY PACK 100-

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers. For Trikafta, severe hepatic impairment.
Required Medical Information	Diagnosis of covered use, submission of cystic fibrosis (CF) mutation test confirming presence of CFTR gene mutations as indicated (see Other Criteria).
Age Restrictions	
Prescriber Restrictions	Restricted to pulmonology
Coverage Duration	1 year
Other Criteria	Initial authorization requires CF mutation test confirming presence of CFTR gene mutations as follows, by drug being requested: (a) for Kalydeco, a mutation predicted to be responsive to ivacaftor based on section 12.1 of the prescribing information, (b) for Orkambi, two copies of the F508del mutation, (c) for Symdeko, two copies of the F508del mutation or at least one mutation predicted to be responsive based on section 12.1 of the prescribing information, (d) for Trikafta, at least one mutation predicted to be responsive based on section 12.1 of the prescribing information or a responsive mutation based on in vitro data.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CHENODAL (chenodiol)

Products Affected

• CHENODAL

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, known hepatocyte dysfunction, bile duct abnormalities such as intrahepatic cholestasis, primary biliary cirrhosis, or sclerosing cholangitis, radiopaque stones, nonvisualizing gallbladder confirmed as nonvisualizing after 2 consecutive single doses of dye, compelling reasons for gallbladder surgery
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to gastroenterology and hepatology
Coverage Duration	24 months
Other Criteria	Safety beyond 24 months is not established and will not be authorized.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CHOLBAM (cholic acid)

Products Affected

• CHOLBAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of liver function testing.
Age Restrictions	
Prescriber Restrictions	Restricted to gastroenterology, hepatology, and pediatric gastroenterology
Coverage Duration	Initially 3 months, then 1 year
Other Criteria	For the first reauthorization, documentation of liver function improvement without complete biliary obstruction or persistent clinical or laboratory indications of worsening liver function or cholestasis is required. For each annual reauthorization, updated liver function testing since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

COMETRIQ (cabozantinib)

Products Affected

- COMETRIQ (100 MG DAILY DOSE) ORAL KIT COMETRIQ (60 MG DAILY DOSE) 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment (Child-Pugh class C), uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of baseline blood pressure reading, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

COPIKTRA (duvelisib)

Products Affected

• COPIKTRA ORAL CAPSULE 15 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Request as first- or second-line therapy, coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of at least two prior systemic therapies tried and failed, submission of pregnancy status for female patients of childbearing potential, attestation patient will receive prophylaxis for Pneumocystis jirovecii pneumonia (PJP) and, if necessary, cytomegalovirus.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CORTICOTROPIN

Products Affected

- CORTROPHIN
- CORTROPHIN GEL

PA Criteria	Criteria Details
Exclusion Criteria	Request for IV administration, treatment of patients under 2 years of age in whom congenital infections are suspected, patients with scleroderma, osteoporosis, systemic fungal infections, ocular herpes simplex, recent surgery, a history of or presence of a peptic ulcer, congestive heart failure, uncontrolled hypertension, primary adrenocortical insufficiency, adrenocortical hyperfunction, or sensitivity to proteins of porcine origin
Required Medical Information	Diagnosis of covered use, submission of blood pressure reading and baseline serum sodium and potassium levels.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 weeks
Other Criteria	For each reauthorization, updated blood pressure, sodium, and potassium levels since the previous authorization are required. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

COTELLIC/ZELBORAF (cobimetinib/vemurafenib)

Products Affected

- COTELLIC
- ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	For cobimetinib, coadministration with moderate or strong CYP3A inhibitors or inducers. For vemurafenib, electrolyte abnormalities that are not correctable, long QT syndrome, coadministration with drugs that prolong the QT interval.
Required Medical Information	Diagnosis of covered use including verification of BRAF V600 mutation as needed for diagnosis, submission of pregnancy status for female patients of childbearing potential. For patients using cobimetinib, submission of left ventricular ejection fraction (LVEF) with a requirement the baseline LVEF is greater than or equal to 50%. For patients using vemurafenib, submission of QTc interval with a requirement the QT interval is less than or equal to 500 msec.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CRESEMBA (isavuconazonium)

Products Affected

• CRESEMBA ORAL CAPSULE 186 MG

PA Criteria	Criteria Details
Exclusion Criteria	Familial short QT syndrome, coadministration with strong CYP3A inhibitors or inducers
Required Medical Information	Diagnosis of covered use.
Age Restrictions	6 years of age or older
Prescriber Restrictions	Restricted to hematology, infectious diseases, and oncology
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

CYSTEAMINE EYE DROPS

Products Affected

- CYSTADROPS
- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, confirmation of corneal cysteine crystal deposits as seen on slit-lamp examination.
Age Restrictions	
Prescriber Restrictions	Restricted to metabolic diseases specialty, ophthalmology, and optometry
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

DALFAMPRIDINE

Products Affected

• dalfampridine er

PA Criteria	Criteria Details
Exclusion Criteria	History of seizure, moderate or severe renal impairment (CrCl less than or equal to 50 mL/min)
Required Medical Information	Diagnosis of covered use, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), confirmation that patient is able to walk.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated creatinine clearance since the previous authorization and confirmation patient is able to walk is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

DASATINIB

Products Affected

 dasanib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg

PA Criteria	Criteria Details
Exclusion Criteria	Uncorrected hypokalemia or hypomagnesemia, coadministration with proton pump inhibitors or H2 receptor antagonists
Required Medical Information	Diagnosis of covered use, submission of serum potassium and magnesium, pregnancy status for female patients of childbearing potential. For adults with resistance or intolerance to prior therapy, documentation of prior therapy.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

DAURISMO (glasdegib)

Products Affected

• DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, confirmation patient will also be receiving cytarabine as part of chemotherapeutic regimen. If patient is under 75 years of age, documentation of comorbidities that preclude use of intensive induction chemotherapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

DEFERASIROX

Products Affected

• deferasirox oral tablet soluble

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment, estimated glomerular filtration rate (eGFR) less than 40 mL/min, platelet count below 50 x 10^9/L, high-risk myelodysplastic syndromes, advanced malignancies
Required Medical Information	Diagnosis of covered use, submission of complete blood count (CBC), liver function testing (LFT), ferritin, and eGFR from the previous 3 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	For each reauthorization, updated ferritin level and platelet count drawn within last 3 months and updated CBC, LFT, and eGFR drawn within the previous 6 months is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

DEFERIPRONE

Products Affected

• deferiprone

PA Criteria	Criteria Details
Exclusion Criteria	Absolute neutrophil count (ANC) below 1.5 x 10^9/L, transfusional iron overload in myelodysplastic syndrome or Diamond Blackfan anemia
Required Medical Information	Diagnosis of covered use, submission of serum ferritin levels, ANC, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated ferritin level and ANC within last 3 months is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

DIACOMIT (stiripentol)

Products Affected

• DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	Requests for monotherapy, moderate or severe renal impairment, moderate or severe hepatic impairment
Required Medical Information	Diagnosis of covered use, confirmation patient is also receiving clobazam.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

DICLOFENAC PATCH

Products Affected

• diclofenac epolamine external

PA Criteria	Criteria Details
Exclusion Criteria	Treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery, use on non-intact or damaged skin resulting from any etiology including exudative dermatitis, eczema, infection lesions, burns, or wounds, pregnancy after 30 weeks gestation
Required Medical Information	Diagnosis of acute pain, defined as short-term pain not lasting longer than a 3-month period.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

DRONABINOL

Products Affected

dronabinol

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. If authorization is requested for treatment of nausea and vomiting associated with cancer therapy, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For initial authorization for treatment of nausea and vomiting associated with cancer therapy, the patient must have tried and failed to have an adequate response to or had an intolerance to at least one 5-HT3 receptor antagonist (e.g., granisetron, ondansetron). If the medication is being administered related to cancer treatment and is a full replacement for intravenous administration of antiemetic therapy within 48 hours of cancer treatment, it is covered as a Part B benefit. To be eligible for Part B coverage, the prescribing physician must indicate this on the prescription.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

ENSACOVE (ensartinib)

Products Affected

• ENSACOVE

PA Criteria	Criteria Details
Exclusion Criteria	Prior ALK-inhibitor use, severe hepatic impairment, coadministration with moderate or strong CYP3A4 inhibitors or inducers or P-glycoprotein inhibitors
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, attestation patient has not previously received an ALK-inhibitor, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

EOHILIA (budesonide oral suspension)

Products Affected

• EOHILIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation of upper endoscopy with biopsy showing at least 15 eosinophils per high-power field or 60 eosinophils/mm2, documentation of signs/symptoms, including but not limited to trouble swallowing, food sticking in esophagus, acid reflux, abdominal or chest pain, or nausea and vomiting, documentation patient has tried and failed at least an 8-week course of proton pump inhibitor therapy (i.e., patient has eosinophilic esophagitis unrelated to gastroesophageal reflux).
Age Restrictions	11 years of age or older
Prescriber Restrictions	Restricted to allergy, gastroenterology, immunology, and otolaryngology/otorhinolaryngology
Coverage Duration	12 weeks
Other Criteria	A maximum of one 12-week course will be allowed every 365 days.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

EPIDIOLEX (cannabidiol)

Products Affected

• EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ERIVEDGE (vismodegib)

Products Affected

• ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

EVRYSDI (risdiplam)

Products Affected

• EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use confirmed by genetic testing including either (a) homozygous deletion of SMN1 exon 7 or (b) compound heterozygosity for SMN1 exon 7 deletion and small mutation, documentation of two or more copies of the SMN2 gene by genetic testing.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For first reauthorization, maintenance of or improvement in any motor score or function compared to baseline is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

FINTEPLA (fenfluramine)

Products Affected

• FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	Administration of monoamine oxidase inhibitors within 14 days of initiation
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

GATTEX (teduglutide)

Products Affected

• GATTEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including confirmation of dependency on parenteral nutrition at least 3 times per week.
Age Restrictions	
Prescriber Restrictions	Restricted to gastroenterology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For the first reauthorization for adults 18 years of age or older, submission of reduction in weekly parenteral nutrition/intravenous support volume from baseline and documentation that a colonoscopy or alternate imaging of the entire colon and upper GI endoscopy with polyp removal and showing no active gastrointestinal malignancy is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

GAVRETO (pralsetinib)

Products Affected

• GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inhibitors, uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of RET gene fusion or mutation, baseline blood pressure reading, pregnancy status for female patients of childbearing potential. For thyroid cancer, attestation patient is radioactive iodine-refractory or ineligible.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

GEFITINIB

Products Affected

• gefinib

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of EGFR exon 19 deletions or exon 21 (L858R) substitution mutations, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

GLP-1 AGONISTS

Products Affected

- MOUNJARO SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.5ML, 12.5 MG/0.5ML, 7.5 MG/0.5ML
- OZEMPIC (0.25 OR 0.5 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 2 • TRULICITY SUBCUTANEOUS SOLUTION MG/3ML
- OZEMPIC (1 MG/DOSE) SUBCUTANEOUS

- SOLUTION PEN-INJECTOR 4 MG/3ML
- OZEMPIC (2 MG/DOSE)
- MG/0.5ML, 15 MG/0.5ML, 2.5 MG/0.5ML, 5 RYBELSUS (FORMULATION R2) ORAL TABLET 1.5 MG, 4 MG, 9 MG
 - RYBELSUS ORAL TABLET 14 MG, 3 MG, 7 MG
 - **AUTO-INJECTOR**

PA Criteria	Criteria Details
Exclusion Criteria	Off-label use for weight management (see Other Criteria)
Required Medical Information	Diagnosis of type 2 diabetes confirmed through one of the following: (1) medical record, or (2) ICD-10 on medical claims, or (3) laboratory results (verifying a hemoglobin A1c greater than or equal to 6.5%, a fasting plasma glucose greater than or equal to 126 mg/dL, a 2-hour postprandial blood glucose greater than or equal to 200 mg/dL after an oral glucose tolerance test, or a random plasma blood glucose greater than or equal to 200 mg/dL combined with classic signs/symptoms of hyperglycemia or hyperglycemic crisis), attestation patient is not receiving another GLP-1 agonist for the treatment of any condition.
Age Restrictions	Age must be consistent with the prescribing information of the drug and condition being treated
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	These products will not be approved for weight management as this off-label use is currently excluded from coverage under Medicare Part D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

GOMEKLI (mirdametinib)

Products Affected

• GOMEKLI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

HEREDITARY ANGIOEDEMA THERAPIES, ACUTE

Products Affected

- icabant acetate subcutaneous soluon prefilled syringe
- sajazir subcutaneous soluon prefilled syringe

PA Criteria	Criteria Details
Exclusion Criteria	Requests for prophylactic hereditary angioedema therapy.
Required Medical Information	Diagnosis of covered use.
Age Restrictions	Age must be consistent with the prescribing information of the drug
Prescriber Restrictions	Restricted to allergy, dermatology, hematology, and immunology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

HEREDITARY ANGIOEDEMA THERAPIES, MAINTENANCE

Products Affected

- HAEGARDA
- TAKHZYRO SUBCUTANEOUS SOLUTION
- TAKHZYRO SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 300 MG/2ML

PA Criteria	Criteria Details
Exclusion Criteria	Requests for acute hereditary angioedema (HAE) therapy (attacks)
Required Medical Information	Diagnosis of covered use, submission of objective or subjective documentation that prophylactic therapy is medically necessary, including, but not limited to activity of disease and disease burden, the frequency of HAE attacks, and quality of life.
Age Restrictions	Age must be consistent with the prescribing information of the drug
Prescriber Restrictions	Restricted to allergy, dermatology, hematology, and immunology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

HERNEXEOS (zongertinib)

Products Affected

HERNEXEOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including documentation the tumor is nonsquamous, submission of test confirming tumor is HER2-positive, pregnancy status for female patients of childbearing potential, documentation of at least one previous systemic therapy that was tried and failed.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

IBRANCE (palbociclib)

Products Affected

• IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming primary tumor type is HR-positive, HER2-negative, attestation the treatment regimen will include concomitant use of an aromatase inhibitor or fulvestrant, pregnancy status for female patients of childbearing potential. For endocrine-resistant breast cancer, submission of genetic tumor testing confirming primary tumor type is PIK3CA-mutated, attestation patient will be taking palbociclib concurrently with inavolisib.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

IBTROZI (taletrectinib)

Products Affected

• IBTROZI

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inhibitors or inducers, proton pump inhibitors, or H2 receptor antagonists
Required Medical Information	Diagnosis of covered use, submission of test confirming tumor is ROS1-positive, pregnancy status for female patients of childbearing potential, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to either crizotinib or entrectinib or have contraindications to both crizotinib and entrectinib.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

IDHIFA (enasidenib)

Products Affected

• IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of IDH2 mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

IMKELDI (imatinib)

Products Affected

• IMKELDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of clinical rationale or documentation detailing why the patient cannot use imatinib oral tablets.
Age Restrictions	
Prescriber Restrictions	Restricted to allergy, dermatology, hematology, and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

IMMUNE GLOBULIN

Products Affected

- GAMMAGARD
- GAMMAGARD S/D LESS IGA
- OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 10 GM/100ML, 10 GM/200ML, 2

GM/20ML, 2.5 GM/50ML, 20 GM/200ML, 5 GM/100ML, 5 GM/50ML

PRIVIGEN

PA Criteria	Criteria Details
Exclusion Criteria	IgA-deficient patients with antibodies against IgA and a history of hypersensitivity.
Required Medical Information	Diagnosis of covered use. For ITP, submission of platelet count. For CLL, documentation of IgG level less than 600 mg/dL and recent history of serious bacterial infection requiring either oral or IV antibiotic therapy. For CIDP, unequivocal diagnosis and documentation patient is refractory, intolerant, or has a contraindication to systemic corticosteroids at therapeutic doses over at least 3 months. For passive immunization against varicella, confirmation that the patient is immunosuppressed and cannot receive varicella-zoster immune globulin.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	For acute conditions/new starts, 3 months. For renewal of chronic conditions, 1 year.
Other Criteria	For each reauthorization for any diagnosis, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required. For IV formulations, covered as a Part B benefit if administered in the home for the treatment of primary immune deficiency. For any other combination of treatment site and indication, additional information may need to be submitted to determine if the immune globulin will be covered as a Part B or Part D benefit.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

INQOVI (decitabine/cedazuridine)

Products Affected

INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

INREBIC (fedratinib)

Products Affected

• INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment, thiamine deficiency, coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of thiamine level (see Other Criteria) and baseline platelet count, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to ruxolitinib. If baseline thiamine level is low, coverage will be delayed until thiamine is repleted.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

INVEGA INJECTABLE (paliperidone injectable suspension)

Products Affected

• INVEGA HAFYERA

MG/1.75ML, 819 MG/2.63ML

 INVEGA TRINZA INTRAMUSCULAR SUSPENSION PREFILLED SYRINGE 273 MG/0.88ML, 410 MG/1.32ML, 546

PA Criteria	Criteria Details
Exclusion Criteria	Dementia-related psychosis
Required Medical Information	Diagnosis of covered use. For the 3-month injection, documentation of at least 4 months' treatment with 1-month paliperidone palmitate extended-release injectable suspension. For the 6-month injection, documentation of at least 4 months' treatment with 1-month paliperidone palmitate extended-release injectable suspension or at least one 3-month injection of 3-month paliperidone palmitate extended-release injectable suspension.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

ITOVEBI (inavolisib)

Products Affected

• ITOVEBI ORAL TABLET 3 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming primary tumor type is HR-positive, HER2-negative, and PIK3CA-mutated, submission of current or previous therapies used to treat the condition (see Other Criteria), attestation that patient has locally advanced or metastatic disease, has not experienced disease progression on or following other PI3K inhibitors, and will be taking inavolisib concurrently with fulvestrant and palbociclib, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to endocrine therapy (e.g., tamoxifen or an aromatase inhibitor) and must not have received prior chemotherapy for metastatic breast cancer. In addition, documentation of clinical rationale why abemaciclib, palbociclib, or ribociclib combined with endocrine therapy is not suitable for the patient must be submitted.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

JAYPIRCA (pirtobrutinib)

Products Affected

JAYPIRCA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria), pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	For initial authorization for mantle cell lymphoma, the patient must have tried and failed to have an adequate response to two previous lines of therapy, including a Bruton's tyrosine kinase (BTK) inhibitor. For initial authorization for chronic lymphocytic leukemia or small lymphocytic lymphoma, the patient must have tried and failed to have an adequate response to two previous lines of therapy, including a BTK inhibitor and a B-cell lymphoma 2 (BCL-2) inhibitor.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

JOENJA (leniolisib)

Products Affected

• JOENJA

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, moderate or severe hepatic impairment (Child-Pugh class B or C)
Required Medical Information	Diagnosis of covered use including submission of test confirming presence of a pathogenic variant of either PIK3CD or PIK3R1, submission of liver function testing or Child-Pugh score, confirmation of negative pregnancy status for female patients of childbearing potential or attestation from physician patient is not pregnant and will be using a highly effective method of contraception, attestation patient is not currently using an immunosuppressive medication.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to specialists in genetic diseases or inborn errors of metabolism
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a clinical benefit (e.g., normalization of lymphocyte subsets, normalization of lymphadenopathy, reduction in spleen size, etc.) or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

KETOCONAZOLE ORAL

Products Affected

ketoconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	Acute or chronic liver disease, treatment of fungal meningitis or fungal infections of the skin or nails
Required Medical Information	Diagnosis of culture-proven systemic blastomycosis, coccidioidomycosis, histoplasmosis, chromomycosis, or paracoccidioidomycosis, submission of baseline ALT, AST, total bilirubin, alkaline phosphatase, prothrombin time and INR, prescriber attestation that the potential benefits of therapy outweigh the risks.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

KISQALI (ribociclib)

Products Affected

- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome, QTcF interval greater than or equal to 450 msec at treatment initiation, uncorrected hypokalemia or hypomagnesemia, coadministration with strong CYP3A4 inducers or drugs that can prolong the QT interval
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming the primary tumor type is HR-positive, HER2-negative, submission of QTcF interval, serum potassium and magnesium drawn within the previous 6 months, pregnancy status for female patients of childbearing potential, attestation that the treatment regimen will include concomitant use of an aromatase inhibitor or fulvestrant.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

KOSELUGO (selumetinib)

Products Affected

KOSELUGO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	Initiation: 2-17 years of age. Continuation: 2 years of age or older.
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	Selumetinib is indicated in pediatric patients and will not be approved for adults unless the patient started on the medication prior to 18 years of age.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

LAZCLUZE (lazertinib)

Products Affected

• LAZCLUZE ORAL TABLET 240 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations, attestation that the medication will be used in combination with amivantamab and will be given with anticoagulant prophylaxis for the first four months of therapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

LEUKINE (sargramostim, GM-CSF)

Products Affected

• LEUKINE INJECTION SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

L-GLUTAMINE

Products Affected

• I-glutamine oral packet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of sickle cell disease, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	
Prescriber Restrictions	Restricted to hematology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must be using, tried and failed to have an adequate response to, or had an intolerance/contraindication to hydroxyurea.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

LIDOCAINE TRANSDERMAL PATCHES

Products Affected

- lidocaine external patch 5 %
- lidocan
- tridacaine ii

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

LIVTENCITY (maribavir)

Products Affected

• LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including a documented history of hematopoietic stem cell or solid organ transplant, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to hematology, infectious diseases, oncology, and transplant specialty
Coverage Duration	8 weeks
Other Criteria	For authorization, the patient must have tried and failed to have an adequate response to at least one of cidofovir, foscarnet, ganciclovir, or valganciclovir.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

LODOCO (colchicine)

Products Affected

• LODOCO

PA Criteria	Criteria Details
Exclusion Criteria	Requests for the treatment of gout, renal failure, severe hepatic impairment, pre-existing blood dyscrasias, coadministration with strong CYP3A4 or P-glycoprotein inhibitors
Required Medical Information	Diagnosis, documented by either (1) prior acute coronary syndrome, or (2) prior ischemic stroke, transient ischemic attack, or carotid artery stenosis greater than 50%, or (3) prior coronary revascularization, or (4) proven coronary artery disease on invasive coronary angiography or computer tomography angiography, or (5) coronary-artery calcium score greater than or equal to 300 Agatston units, or (6) aortic atherosclerotic disease, or (7) symptomatic peripheral vascular disease, submission of estimated glomerular filtration rate (eGFR) or creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance) with a requirement the eGFR or creatinine clearance is greater than or equal to 15 mL/min, and attestations patient (1) does not have severe hepatic impairment, and (2) has had a recent complete blood count and does not have evidence of any cytopenia, and (3) does not have NYHA functional Class 3 or 4 heart failure.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, submission of updated eGFR or creatinine clearance and complete blood count since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
10/1/2025	-

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	No

LORBRENA (lorlatinib)

Products Affected

• LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers, uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, baseline blood pressure, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

LYBALVI (olanzapine/samidorphan)

Products Affected

LYBALVI

PA Criteria	Criteria Details
Exclusion Criteria	Dementia-related psychosis, coadministration with opioids, levodopa, dopamine agonists, or strong CYP3A inducers, acute opioid withdrawal, end-stage renal disease
Required Medical Information	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to neurology and psychiatry
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to both (1) generic olanzapine, including documentation showing a positive therapeutic benefit but unacceptable weight gain as a result of the drug, and (2) one other generic second-generation antipsychotic with low incidence of metabolic side effects (e.g., aripiprazole, lurasidone, ziprasidone).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

MAVYRET (glecaprevir/pibrentasvir)

Products Affected

MAVYRET

PA Criteria	Criteria Details
Exclusion Criteria	Moderate or severe hepatic impairment (Child-Pugh class B or C), coadministration with rifampin or atazanavir
Required Medical Information	Diagnosis of covered use, laboratory confirmation of hepatitis C virus (HCV) infection and HCV genotype.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Criteria for coverage duration will be applied consistent with current AASLD-IDSA guidance.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

MEGESTROL IN OLDER PATIENTS

Products Affected

 megestrol acetate oral suspension 40 mg/ml, 625 mg/5ml

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	PA not required for hematology or oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

MEKINIST/TAFINLAR (trametinib/dabrafenib)

Products Affected

- MEKINIST
- TAFINLAR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of BRAF V600E or V600K mutation, pregnancy status for female patients of childbearing potential. For anaplastic thyroid cancer, BRAF V600E-mutated solid tumors, low-grade glioma, and adjuvant BRAF V600E-and/or V600K-mutated melanoma indications, confirmation that trametinib and dabrafenib will be co-administered.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

MEMANTINE/DONEPEZIL

Products Affected

- memanne hcl-donepezil hcl
- NAMZARIC ORAL CAPSULE EXTENDED RELEASE 24 HOUR 7-10 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of documentation that the patient has tolerated donepezil 10 mg daily for a minimum of 1 month.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

MODEYSO (dordaviprone)

Products Affected

MODEYSO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming an H3K27M mutation, pregnancy status for female patients of childbearing potential, documentation of at least one previous therapy.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

NILOTINIB

Products Affected

- DANZITEN
- nilonib hcl

PA Criteria	Criteria Details
Exclusion Criteria	Uncorrected hypokalemia or hypomagnesemia, long QT syndrome, coadministration with drugs that prolong the QT interval, proton pump inhibitors, or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, confirmation of positive Philadelphia chromosome (Ph) status, baseline serum potassium and magnesium levels.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

NINLARO (ixazomib)

Products Affected

NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, documentation that medication will be administered concomitantly with lenalidomide and dexamethasone, documentation of prior therapy regimen for multiple myeloma, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

NITISINONE

Products Affected

• nisinone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of succinylacetone in urine or plasma.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a clinical benefit, such as reductions in urine succinylacetone level, alphafetoprotein level, serum tyrosine level, or serum phenylalanine level, or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

NUCALA (mepolizumab)

Products Affected

- NUCALA SUBCUTANEOUS SOLUTION AUTO NUCALA SUBCUTANEOUS SOLUTION **INJECTOR**
- NUCALA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 100 MG/ML
- RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	Regular (non-eosinophilic) granulomatosis with polyangiitis (GPA) or microscopic polyangiitis (MPA)
Required Medical Information	Diagnosis of covered use. For asthma, chronic rhinosinusitis with nasal polyps (CRSwNP), and hypereosinophilic syndrome (HES), submission of current and previous therapies used to treat the condition (see Other Criteria). For asthma, (1) documentation of a pre-bronchodilator FEV1 less than 80% predicted in adults, or less than 90% in children, and (2) submission of blood eosinophil (eos) count of at least 150 cells/mcL obtained within 6 weeks of therapy initiation or at least 300 cells/mcL within 12 months of therapy initiation. For CRSwNP, (1) documentation of evidence of nasal polyps, and (2) attestation that patient has symptomatic nasal congestion. For eosinophilic granulomatosis with polyangiitis, documentation of (1) a history of asthma, and (2) an eos percentage greater than or equal to 10% or an absolute eos count greater than 1000 cells/cubic millimeter from the previous 6 weeks, and (3) disease lasting at least 6 months that is relapsed or refractory to oral corticosteroids and/or immunosuppressive therapies. For HES, documentation of (1) uncontrolled disease defined as a history of at least 2 flares requiring systemic therapy within the past 12 months and a blood eos count of at least 1000 cells/mcL from the previous 6 weeks, and (2) attestation disease does not have an identifiable non-hematologic secondary cause.
Age Restrictions	
Prescriber Restrictions	Restricted to allergy, hematology, immunology, otorhinolaryngology, pulmonology, and rheumatology
Coverage Duration	1 year

PA Criteria	Criteria Details
Other Criteria	For initial authorization for asthma, patient must be on a drug regimen as recommended by GINA guidelines prior to the use of biologic medications, consisting of, at a minimum, a maximally-tolerated dose of inhaled corticosteroid (ICS), a long-acting beta-agonist (LABA), and a long-acting antimuscarinic antagonist (LAMA), and the provider must attest this therapy will be continued after starting mepolizumab. For initial authorization for CRSwNP, patient must have tried an intranasal corticosteroid for at least two months (and provider must attest this will be continued after starting mepolizumab), have a contraindication to intranasal corticosteroids, or documentation must be submitted as to why this therapy is not otherwise advisable. For initial authorization for HES, patient must have been using a course of oral corticosteroids, cytotoxic therapy, or immunosuppressive therapy for at least the previous 4 weeks. For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required. For each reauthorization for asthma, confirmation patient is still using triple ICS-LABA-LAMA inhaler therapy is required. For each reauthorization for CRSwNP, confirmation patient is still using an intranasal corticosteroid, unless contraindicated, is required. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

NUPLAZID (pimavanserin)

Products Affected

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

PA Criteria	Criteria Details
Exclusion Criteria	Psychosis unrelated to Parkinson's disease psychosis, cardiac arrhythmias, symptomatic bradycardia, congenital QT prolongation, coadministration with moderate or strong CYP3A4 inducers or drugs that prolong the QT interval, uncorrected hypokalemia or hypomagnesemia
Required Medical Information	Diagnosis of covered use, submission of baseline serum potassium and magnesium levels.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ODOMZO (sonidegib)

Products Affected

• ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, coadministration with strong CYP3A4 inhibitors or moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, attestation patient is not a candidate for surgery or radiation therapy or carcinoma has recurred following surgery or radiation therapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

OGSIVEO (nirogacestat)

Products Affected

• OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inhibitors or inducers, proton pump inhibitors, or H2 receptor antagonists
Required Medical Information	Diagnosis of covered use with documentation of tumor progression, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology and sarcoma specialty
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

OJEMDA (tovorafenib)

Products Affected

OJEMDA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP2C8 inhibitors or inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of BRAF V600 mutation or BRAF gene fusion or rearrangement, documentation of previous systemic therapy/therapies for pediatric low-grade glioma tried and failed with a minimum of one previous therapy necessary for approval, pregnancy status for female patients of childbearing potential. If genetic testing does not reveal a BRAF gene fusion or rearrangement, documentation of previous intolerance to, contraindication to, or other reason why the patient cannot use the combination of trametinib and dabrafenib.
Age Restrictions	Initiation: 21 years of age or younger (see Other Criteria)
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	Tovorafenib is indicated as therapy in children and young adults and will not be approved for adults unless the patient started on the medication prior to 22 years of age.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

OJJAARA (momelotinib)

Products Affected

• OJJAARA ORAL TABLET 100 MG, 150 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	Active infection, uncontrolled acute or chronic liver disease
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ONUREG (azacitidine)

Products Affected

• ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of myelodysplastic syndrome
Required Medical Information	Diagnosis of covered use, documentation patient achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy and attestation patient cannot complete intensive curative therapy, submission of absolute neutrophil count (with the requirement it is at least 500/mcL), submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ORIAHNN (elagolix/estradiol/norethindrone)

Products Affected

ORIAHNN

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, known liver impairment or disease, known osteoporosis, undiagnosed abnormal uterine bleeding, women who are at increased risk of, have a history of, or currently have thrombotic or thromboembolic disorders (including women over 35 years of age who smoke and women with uncontrolled hypertension), current/history of breast cancer or other hormone-sensitive cancer
Required Medical Information	Diagnosis of covered use, attestation patient is premenopausal, submission of baseline blood pressure, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to endocrinology and gynecology
Coverage Duration	2 years (see Other Criteria)
Other Criteria	Use of this drug for more than 2 years increases risk of bone loss and requests for therapy for more than 2 years will not be approved.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ORILISSA (elagolix)

Products Affected

• ORILISSA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, severe hepatic impairment (Child-Pugh class C), known osteoporosis, coadministration with OATP1B1 inhibitors
Required Medical Information	Diagnosis of covered use, attestation patient is premenopausal and will be using non-hormonal contraception during therapy, submission of liver function testing or Child-Pugh score, pregnancy status for female patients of childbearing potential. If the patient has previously used elagolix, submission of dose used and number of total months of prior use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to endocrinology and gynecology
Coverage Duration	Up to 24 months based on dose and coexisting conditions (see Other Criteria)
Other Criteria	Due to increased risk of bone loss, maximum duration of use is limited based on dose and coexisting conditions. For (1) endometriosis with dyspareunia where dose will be 200 mg twice daily, or (2) women with moderate hepatic impairment, the maximum duration of use is 6 months. Requests for use greater than 6 months will not be approved in these situations. For (3) endometriosis without dyspareunia, 150 mg daily for 24 months. Requests for use greater than 24 months will not be approved in this situation.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

PANRETIN (alitretinoin)

Products Affected

• PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, instances where systemic Kaposi sarcoma therapy is required (more than 10 new Kaposi's sarcoma lesions in the prior month, symptomatic lymphedema, symptomatic pulmonary Kaposi sarcoma, or symptomatic visceral involvement)
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

PAZOPANIB

Products Affected

• pazopanib hcl

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment, uncontrolled hypertension, uncorrected hypokalemia, hypocalcemia, or hypomagnesemia, coadministration with strong CYP3A4 inducers, proton pump inhibitors, H2-receptor antagonists, or drugs that can prolong the QT interval
Required Medical Information	Diagnosis of covered use, submission of baseline blood pressure, serum potassium, calcium, and magnesium levels, pregnancy status for female patients of childbearing potential. For soft tissue sarcoma, submission of previous chemotherapy regimen(s).
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

PEGFILGRASTIM

Products Affected

- UDENYCA
- UDENYCA ONBODY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of FDA-approved indication.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

PEMAZYRE (pemigatinib)

Products Affected

• PEMAZYRE ORAL TABLET 13.5 MG, 4.5 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of either FGFR1 rearrangement or FGFR2 fusion or rearrangement depending on the indication, attestation patient has used previous systemic treatment for the indication, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

PIRFENIDONE

Products Affected

• pirfenidone oral tablet 267 mg, 801 mg

PA Criteria	Criteria Details
Exclusion Criteria	End-stage renal disease on dialysis, severe hepatic impairment (Child- Pugh class C)
Required Medical Information	Diagnosis of covered use, submission of liver function tests or Child-Pugh score.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to pulmonology
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved and updated liver function testing or Child-Pugh score since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

PRETOMANID

Products Affected

PRETOMANID

PA Criteria	Criteria Details
Exclusion Criteria	Inability to use bedaquiline or linezolid, drug-sensitive tuberculosis, coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, attestation pretomanid will be used in combination with bedaquiline and linezolid.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to infectious diseases and pulmonology
Coverage Duration	26 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

PREVYMIS (letermovir)

Products Affected

• PREVYMIS ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment (Child-Pugh class C), coadministration with ergot alkaloids, pimozide, or pitavastatin or simvastatin when coadministered with cyclosporine
Required Medical Information	Diagnosis of covered use, submission of day number post-transplant. For use after kidney transplant, documentation patient is high risk, defined as donor CMV seropositive/recipient CMV seronegative (D+/R-), submission of explanation why valganciclovir is contraindicated or cannot be used for prophylaxis.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology, infectious diseases, oncology, and transplant speciality
Coverage Duration	100 days post-HSCT or 200 days post-kidney transplant or post-HSCT at risk for late CMV infection
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

PROMETHAZINE IN OLDER PATIENTS

Products Affected

- promethazine hcl oral soluon 6.25 mg/5ml
- promethazine hcl oral tablet
- promethazine hcl rectal suppository 12.5 mg,
 25 mg
- PROMETHEGAN RECTAL SUPPOSITORY 25 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. For allergic conditions, documentation must be submitted showing patient has tried and failed or had an inadequate response to a second-generation antihistamine.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	Promethazine is a potent anticholinergic considered high-risk in older patients due to risks of confusion, dry mouth, constipation, and decreased clearance with advanced age.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

PROSTATE CANCER ORAL MEDICATIONS

Products Affected

- AKEEGA
- ERLEADA
- NUBEQA
- XTANDI

PA Criteria	Criteria Details
Exclusion Criteria	For Akeega, severe hepatic impairment (Child-Pugh class C), uncontrolled hypertension, uncontrolled hypokalemia
Required Medical Information	Diagnosis of covered use. For Nubeqa, submission of current or previous therapies used to treat the condition (see Other Criteria). For Akeega, submission of test confirming presence of deleterious BRCA mutation, Child-Pugh score or liver function testing, baseline blood pressure reading, and serum potassium level.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology and urology
Coverage Duration	1 year
Other Criteria	For initial authorization of Nubeqa, the patient must have tried and failed to have an adequate response or had an intolerance to both Erleada and Xtandi.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

PULMONARY HYPERTENSION MEDICATIONS

Products Affected

- alyq
- ambrisentan oral tablet 10 mg, 5 mg
- bosentan oral tablet 125 mg, 62.5 mg
- OPSUMIT
- ORENITRAM
- ORENITRAM MONTH 1

- ORENITRAM MONTH 2
- ORENITRAM MONTH 3
- sildenafil citrate oral suspension reconstuted
- sildenafil citrate oral tablet 20 mg
- tadalafil (pah)

PA Criteria	Criteria Details
FA CITICITA	Citteria Details
Exclusion Criteria	For ambrisentan, bosentan, or Opsumit, pregnancy. For ambrisentan or Orenitram, moderate or severe hepatic impairment. For tadalafil, severe hepatic impairment or creatinine clearance below 30 mL/min or on hemodialysis. For ambrisentan, idiopathic pulmonary fibrosis.
Required Medical Information	Diagnosis of covered use confirmed by right heart catheterization, submission of mean pulmonary arterial pressure greater than 20 mm Hg at rest, pulmonary arterial wedge pressure less than or equal to 15 mm Hg, pulmonary vascular resistance greater than 2 Woods units, and attestation patient is WHO Group 1. For ambrisentan, bosentan, or Opsumit, submission of pregnancy status for female patients of childbearing potential. For Opsumit only, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	For all drugs in this policy except bosentan, 18 years of age or older
Prescriber Restrictions	Restricted to cardiology and pulmonology
Coverage Duration	1 year
Other Criteria	For initial authorization of Opsumit, the patient must have tried and failed to have an adequate response to or had an intolerance to ambrisentan or bosentan.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

RETEVMO (selpercatinib)

Products Affected

• RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	Uncontrolled hypertension, coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of RET gene fusion or mutation, baseline blood pressure reading, pregnancy status for female patients of childbearing potential. For RET fusion-positive thyroid cancer, attestation patient is radioactive iodine-refractory or ineligible. For solid tumors with a RET gene fusion, documentation of previous systemic therapy tried or reason why patient has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

REVCOVI (elapegademase-lvlr)

Products Affected

REVCOVI

PA Criteria	Criteria Details
Exclusion Criteria	Severe thrombocytopenia
Required Medical Information	Diagnosis of covered use confirmed by either biochemical testing showing less than 1% of adenosine deaminase (ADA) catalytic activity in red blood cells or genetic testing showing biallelic ADA pathogenic variants, submission of baseline plasma ADA level (if genetic testing was submitted as confirmation of diagnosis) and platelet count.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology, immunology, and specialists in genetic diseases
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, submission of updated plasma ADA level demonstrating an improvement from baseline and platelet count is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

REVUFORJ (revumenib)

Products Affected

• REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	QTcF interval greater than 450 msec at treatment initiation, white blood cell (WBC) count greater than 25 x 10^9/L, uncorrected hypokalemia or hypomagnesemia, coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of a lysine methyltransferase 2A gene (KMT2A) translocation that is not a 11q23 partial tandem duplication, submission of baseline QTcF interval, serum potassium, serum magnesium, baseline WBC count, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

REZLIDHIA (olutasidenib)

Products Affected

• REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of IDH1 mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

RIVFLOZA (nedosiran)

Products Affected

• RIVFLOZA

PA Criteria	Criteria Details
Exclusion Criteria	Estimated glomerular filtration rate (eGFR) less than 30 mL/min/1.73 m2
Required Medical Information	Diagnosis of covered use, documentation of AGXT mutation confirmed by liver enzyme analysis or genetic testing, submission of 24-hour urinary oxalate (Uox) excretion with a requirement it is greater than or equal to 0.7 mmol (normalized to body surface area if patient is under 18 years of age) and eGFR, attestation patient has not received a prior kidney or liver transplant, attestation patient will not be using in combination with lumasiran (Oxlumo).
Age Restrictions	
Prescriber Restrictions	Restricted to nephrology and urology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For the first reauthorization, documentation of clinically relevant response to therapy as evidenced by reduced Uox or plasma oxalate levels is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ROMVIMZA (vimseltinib)

Products Affected

• ROMVIMZA

PA Criteria	Criteria Details
Exclusion Criteria	Active liver or biliary tract disease (including increased ALP), pre-existing increased serum transaminases, total or direct bilirubin greater than the upper limit of normal
Required Medical Information	Diagnosis of covered use (and documentation surgical intervention is not possible or practical), submission of serum transaminases, total and direct bilirubin, and ALP, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ROZLYTREK (entrectinib)

Products Affected

ROZLYTREK

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers or drugs that prolong the QTc interval
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For non-small cell lung cancer, submission of test confirming presence of ROS1-positive tumor. For solid tumors, submission of evidence of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation and attestation tumor is metastatic or surgical resection/other systemic therapies are unsatisfactory treatment options.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

RYDAPT (midostaurin)

Products Affected

RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For acute myeloid leukemia, submission of test confirming presence of FLT3 mutation, attestation patient will be receiving cytarabine and daunorubicin induction and cytarabine consolidation with midostaurin.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to allergy, hematology, and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

SAPROPTERIN

Products Affected

- javygtor
- sapropterin dihydrochloride oral packet
- sapropterin dihydrochloride oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of blood phenylalanine concentration.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For the first reauthorization, documentation of reduction in blood phenylalanine concentration from pre-treatment baseline is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

SECOND-GENERATION ANTIPSYCHOTICS

Products Affected

- CAPLYTA ORAL CAPSULE 10.5 MG, 21 MG, 42 MG
- COBENFY
- COBENFY STARTER PACK
- FANAPT ORAL TABLET 1 MG, 10 MG, 12 MG, SECUADO TRANSDERMAL PATCH 24 HOUR 2 MG, 4 MG, 6 MG, 8 MG
- FANAPT TITRATION PACK A

- FANAPT TITRATION PACK B ORAL TABLET
- FANAPT TITRATION PACK C ORAL TABLET
- REXULTI ORAL TABLET 0.25 MG, 0.5 MG, 1 MG, 2 MG, 3 MG, 4 MG
- 3.8 MG/24HR, 5.7 MG/24HR, 7.6 MG/24HR

PA Criteria	Criteria Details
Exclusion Criteria	Dementia-related psychosis. For Cobenfy only, moderate to severe renal impairment, urinary retention, gastric retention, hepatic impairment, untreated narrow-angle glaucoma.
Required Medical Information	Diagnosis of covered use. For schizophrenia, an indication related to bipolar disorder type I, or for Rexulti for major depressive disorder, submission of current or previous therapies used to treat the condition (see Other Criteria). For Cobenfy only, submission of estimated glomerular filtration rate with the requirement it is at least 60 mL/min, attestation patient does not have hepatic impairment.
Age Restrictions	
Prescriber Restrictions	Restricted to psychiatry
Coverage Duration	1 year

PA Criteria	Criteria Details
Other Criteria	For initial authorization for schizophrenia or acute treatment of manic/mixed episodes of bipolar I disorder, the patient must have tried and failed to have an adequate response to or had an intolerance to aripiprazole and at least one other generic second-generation atypical antipsychotic. For initial authorization of Caplyta for depressive episodes associated with bipolar I disorder, the patient must have tried and failed to have an adequate response to or had an intolerance to at least two of the following drugs: cariprazine, lurasidone, olanzapine, or quetiapine. For Rexulti as an adjunctive therapy to antidepressants for major depressive disorder, the patient must have tried and failed to have an adequate response to or had an intolerance to aripiprazole and cariprazine or quetiapine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

SEDATIVE HYPNOTICS IN OLDER PATIENTS

Products Affected

- eszopiclone
- zaleplon
- zolpidem tartrate oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation at least two of the following medications were tried and deemed ineffective or intolerable: Belsomra, doxepin tablets, ramelteon, and trazodone.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	Sedative hypnotic medications are high-risk medications in older patients due to increased risks of cognitive impairment, delirium, unsteady gait, syncope, falls, fractures, and motor vehicle accidents.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

SOFOSBUVIR/VELPATASVIR

Products Affected

• sofosbuvir-velpatasvir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, laboratory confirmation of hepatitis C virus (HCV) infection and HCV genotype, attestation that patients with decompensated cirrhosis will receive concomitant ribavirin therapy unless ribavirin therapy is contraindicated.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

SOHONOS (palovarotene)

Products Affected

SOHONOS

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of R206H ACVR1 mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	For female patients, 8 years of age or older. For male patients, 10 years of age or older.
Prescriber Restrictions	Restricted to orthopedics, rheumatology, and specialists in rare connective tissue diseases
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved and attestation the patient continues to undergo regular pregnancy testing (as necessary for patients of childbearing potential) is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

SOMAVERT (pegvisomant)

Products Affected

• SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including attestation that surgery or radiation was not curative or is not an option, submission of baseline IGF-1, submission of baseline liver function testing (LFT) including bilirubin with the requirement liver transaminases are less than or equal to 3 times the upper limit of normal (ULN). If liver transaminases are greater than 3 times ULN, submission of the cause of liver dysfunction determined through a comprehensive workup.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to endocrinology
Coverage Duration	1 year
Other Criteria	For the first reauthorization, submission of updated IGF-1 level demonstrating an improvement from baseline, LFT showing liver transaminases below 5 times the ULN, and attestation patient does not have signs or symptoms of liver injury (e.g., jaundice, elevated bilirubin level or bilirubinuria, fatigue, nausea, vomiting, right upper quadrant pain, ascites, unexplained edema, easy bruisability) is required. For each annual reauthorization, documented improvement/maintenance of IGF-1 level is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

SORAFENIB

Products Affected

• sorafenib tosylate

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome, coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For differentiated thyroid cancer, attestation patient is radioactive iodine-refractory or ineligible.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

TABRECTA (capmatinib)

Products Affected

• TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of MET exon 14 skipping mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

TAFAMIDIS

Products Affected

- VYNDAMAX
- VYNDAQEL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of transthyretin amyloid cardiomyopathy (ATTRwt or ATTRm) confirmed by one of the following: (1) presence of amyloid deposits on cardiac biopsy, (2) presence of transthyretin precursor protein confirmed on immunohistochemical analysis, (3) technetium 99-labeled nuclear scintigraphy or single-photon emission computer tomography (SPECT), or (4) a TTR genetic mutation plus cardiac involvement defined as thickening of the interseptal ventricular wall. In addition, patients should also have documentation of history of heart failure, with at least one prior hospitalization for heart failure or clinical evidence of heart failure with signs or symptoms of volume overload requiring treatment with a diuretic for improvement.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

TAGRISSO (osimertinib)

Products Affected

• TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of EGFR exon 19 deletions, exon 21 L858R mutations, or T790M mutations, pregnancy status for female patients of childbearing potential. For EGFR T790M mutation-positive NSCLC, submission of previous EGFR tyrosine kinase inhibitor therapy used for indication.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

TALZENNA (talazoparib)

Products Affected

 TALZENNA ORAL CAPSULE 0.1 MG, 0.25 MG, 0.35 MG, 0.5 MG, 0.75 MG, 1 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For breast cancer, submission of test results confirming germline BRCA mutation-positive, human epidermal growth factor receptor 2 (HER2) negative disease. For prostate cancer, submission of test results confirming HRR gene-mutated disease, confirmation talazoparib will be used in combination with enzalutamide.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

TAVNEOS (avacopan)

Products Affected

TAVNEOS

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers, active serious infection, chronic active hepatitis B, untreated hepatitis C, uncontrolled autoimmune hepatitis, cirrhosis
Required Medical Information	Diagnosis of covered use (granulomatosis with polyangiitis or microscopic polyangiitis variants of anti-neutrophil cytoplasmic antibody [ANCA]-associated vasculitis), submission of HBV serology testing, attestation patient is using rituximab, cyclophosphamide/azathioprine, or another compendium-supported therapy for the treatment of ANCA-associated vasculitis, along with glucocorticoids.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to immunology, nephrology, pulmonology, and rheumatology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For the first reauthorization, documentation of clinically relevant response to therapy, including but not limited to disease remission defined using changes in Birmingham Vasculitis Activity Score, a documented reduction in maintenance glucocorticoid dose, or improved or sustained renal function, is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

TAZVERIK (tazemetostat)

Products Affected

TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For relapsed/refractory follicular lymphoma, documentation (1) of test confirming presence of EZH2 mutation and treatment with at least two prior systemic therapies, or (2) patient has no satisfactory alternative treatment option.
Age Restrictions	16 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

TEPMETKO (tepotinib)

Products Affected

• TEPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of MET exon 14 skipping mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

TERIPARATIDE

Products Affected

• teriparade subcutaneous soluon peninjector 560 mcg/2.24ml

PA Criteria	Criteria Details
Exclusion Criteria	Pre-existing hypercalcemia, underlying hypercalcemic disorder (such as primary hyperparathyroidism), patients with an increased risk of osteosarcoma (such as those with Paget's disease)
Required Medical Information	Diagnosis of covered use where "high risk for fracture" is defined as documentation of (1) a history of fracture of the hip or vertebra regardless of bone mineral density (BMD), or (2) a history of fracture of the proximal humerus, pelvis, or distal forearm and T-score between -1.0 and -2.5, or (3) T-score less than or equal to -2.5 at the total hip, femoral neck, spine, or distal third of the radius, or (4) T-score between -1.0 and -2.5 at the total hip, femoral neck, spine, or distal third of the radius, and (a) a 10-year probability of hip fracture as assessed by FRAX score of at least 3%, or (b) a 10-year probability of a major osteoporosis-related fracture as assessed by FRAX score of at least 20%, submission of baseline serum calcium, postmenopausal status, current or previous therapies used to treat the condition (see Other Criteria), number of total months of all prior use of parathyroid hormone analogs and parathyroid hormone related peptides.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	2 years unless patient is at high risk for fracture after 2 years of therapy (see Other Criteria)

PA Criteria	Criteria Details
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to at least one bisphosphonate. Therapeutic failure is defined as either a fracture or a decrease in BMD while using a bisphosphonate for at least 3 months. Use of parathyroid hormone analogs and/or parathyroid hormone related peptides for more than 2 years during a patient's lifetime is generally not recommended. For annual reauthorization beyond 2 years, submission of updated serum calcium since initial authorization and evidence the patient remains at high risk for fracture as defined in the Required Medical Information section is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

TOLVAPTAN (HYPONATREMIA)

Products Affected

 tolvaptan oral tablet 15 mgtolvaptan (hyponatremia), 30 mgtolvaptan (hyponatremia)

PA Criteria	Criteria Details
Exclusion Criteria	Underlying liver disease, need to raise serum sodium acutely, inability to sense or respond to thirst, hypovolemia, anuria, coadministration with strong CYP3A inhibitors or inducers or desmopressin
Required Medical Information	Diagnosis of covered use, submission of evidence of clinically significant hyponatremia, defined as (1) serum sodium less than 125 mEq/L, or (2) serum sodium less than 135 mEq/L that is symptomatic and has resisted correction with fluid restriction.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	30 days (see Other Criteria)
Other Criteria	Treatment should be initiated in a setting where serum sodium can be monitored closely. Treatment is limited to 30 days to prevent liver injury. This formulation of tolvaptan will not be approved for autosomal dominant polycystic kidney disease (ADPKD) as it is not indicated for ADPKD and the tolvaptan formulation approved for ADPKD has a mandatory REMS program, making it only available through a restricted distribution program.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

TURALIO (pexidartinib)

Products Affected

• TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	Active liver or biliary tract disease (including increased ALP), pre-existing increased serum transaminases, total or direct bilirubin greater than the upper limit of normal, coadministration with other hepatotoxic medications, strong CYP3A inducers, or proton pump inhibitors
Required Medical Information	Diagnosis of covered use (and documentation surgical intervention is not possible or practical), documentation of patient's severe morbidity or functional limitations, submission of serum transaminases, total and direct bilirubin, and ALP, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VALCHLOR (mechlorethamine)

Products Affected

VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	Use as initial therapy
Required Medical Information	Diagnosis of covered use, submission of previous skin-directed therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VANFLYTA (quizartinib)

Products Affected

VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	Requests for maintenance monotherapy after allogeneic hematopoietic stem cell transplant, uncorrected hypokalemia or hypomagnesemia, long QT syndrome, QTcF interval greater than 450 msec at treatment initiation, coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use including submission of test confirming presence of FLT3 internal tandem duplication-positive mutation, submission of QTcF interval, baseline serum potassium and magnesium levels, and pregnancy status for female patients of childbearing potential, attestation patient does not have history of ventricular arrhythmias or torsades de pointes.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VENCLEXTA (venetoclax)

Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers. For CLL/SLL, coadministration with strong CYP3A inhibitors at treatment initiation and initial dosage titration.
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VEOZAH (fezolinetant)

Products Affected

VEOZAH

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with CYP1A2 inhibitors, severe renal impairment or end-stage renal disease, known cirrhosis, any aminotransferase or total bilirubin at or above 2 times the upper limit of normal
Required Medical Information	Diagnosis of covered use, submission of estimated glomerular filtration rate, liver function testing, current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to at least two prior systemic hormone therapies or FDA-approved or compendia-supported non-hormonal therapies (e.g., SSRI, SNRI, clonidine, gabapentin, etc.) for the treatment of vasomotor symptoms due to menopause. The drugs tried must come from different medication classes. For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

VERZENIO (abemaciclib)

Products Affected

• VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers or ketoconazole
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming that the primary tumor type is HR-positive, HER2-negative, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VIJOICE (alpelisib)

Products Affected

- VIJOICE ORAL PACKET
- VIJOICE ORAL TABLET THERAPY PACK 125 MG, 200 & 50 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use including at least one target lesion on imaging with requesting provider attestation patient has severe or lifethreatening disease, submission of test confirming presence of mutation in PIK3CA gene, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to specialists in genetic diseases or inborn errors of metabolism
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a symptomatic or clinical benefit (e.g., reductions in target lesion size, pain, vascular malformations, limb enlargements, etc.), or maintenance of improvement previously achieved, is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VITRAKVI (larotrectinib)

Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of evidence of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation, attestation tumor is metastatic or surgical resection and other systemic therapies are unsatisfactory treatment options, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VIZIMPRO (dacomitinib)

Products Affected

• VIZIMPRO ORAL TABLET 15 MG, 30 MG, 45 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with a proton pump inhibitor
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of EGFR exon 19 deletions or exon 21 L858R substitution mutations, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VONJO (pacritinib)

Products Affected

VONJO

PA Criteria	Criteria Details
Exclusion Criteria	Moderate or severe hepatic impairment (Child-Pugh class B or C), estimated glomerular filtration rate (eGFR) less than 30 mL/min, QTc interval greater than 480 msec at baseline, uncorrected hypokalemia, coadministration with strong CYP3A4 inducers or strong CYP3A4 inhibitors
Required Medical Information	Diagnosis of covered use, submission of platelet count, serum potassium level, eGFR, and QTc interval, Child-Pugh score, documentation from a physical exam patient has splenomegaly.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VOQUEZNA (vonoprazan)

Products Affected

• VOQUEZNA ORAL TABLET 10 MG, 20 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. For erosive esophagitis and non-erosive gastroesophageal reflux disease (GERD), submission of current or previous therapies used to treat the condition (see Other Criteria). For Helicobacter pylori infection only, attestation patient will be administering with amoxicillin or a combination of amoxicillin and clarithromycin.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	Up to 32 weeks based on covered use (see Other Criteria)
Other Criteria	For initial authorization for erosive esophagitis and non-erosive GERD, the patient must have tried and failed to have an adequate response to two different proton pump inhibitors or have contraindications to the proton pump inhibitor class. For non-erosive GERD, the initial coverage duration will be 4 weeks, with the option to reauthorize for an additional 20 weeks (for a total of 24 weeks of therapy per 365 days) if the provider attests to medical need. For H. pylori infection, a maximum of one 14-day course of vonoprazan will be approved. For all other indications, a maximum of one 32-week course of vonoprazan will be allowed per 365 days.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

VORANIGO (vorasidenib)

Products Affected

• VORANIGO ORAL TABLET 10 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of IDH1 or IDH2 mutation, pregnancy status for female patients of childbearing potential, attestation patient has had at least one prior surgery (biopsy, sub-total resection, or gross total resection).
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

VOWST (fecal microbiota spores, live-brpk)

Products Affected

VOWST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use with the requirement patient is being treated after at least 2 recurrent (3 total) Clostridioides difficile infections (confirmation of pathogen with stool test or other confirmatory test), submission of time of last planned dose of antibiotic for latest recurrent C. difficile infection and attestation patient will be using a bowel cleanse the evening prior to starting Vowst, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 course (3 days)
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to or had a contraindication to fecal microbiota, livejslm rectal suspension.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	Yes
Prerequisite Therapy Required	No

WINREVAIR (sotatercept-csrk)

Products Affected

 WINREVAIR SUBCUTANEOUS KIT 2 X 45 MG, 2 X 60 MG, 45 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	Platelet count below 50 x 10^9/L, pregnancy
Required Medical Information	Diagnosis of covered use confirmed by right heart catheterization, submission of mean pulmonary arterial pressure greater than 20 mm Hg at rest, pulmonary arterial wedge pressure less than or equal to 15 mm Hg, pulmonary vascular resistance greater than 2 Woods units, New York Heart Association functional class with the requirement the patient falls into Class II, III, or IV, attestation patient is WHO Group 1, submission of baseline platelet count, current or previous therapies used to treat the condition (see Other Criteria), and pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology and pulmonology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For initial authorization, the patient's drug regimen must be submitted and must include at minimum (1) a nitric oxide pathway mediator, and (2) an endothelin receptor antagonist, and (3) a prostacyclin pathway agonist. If any of these three components are not currently being used, prescriber is required to submit documentation as to why (e.g., intolerances, physiologic parameters, contraindications, etc.). For each reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

XALKORI (crizotinib)

Products Affected

XALKORI

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome, coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming tumor is ALK or ROS1-positive, pregnancy status for female patients of childbearing potential.
Age Restrictions	For ALK-positive systemic anaplastic large cell lymphoma only, 1 year of age to 21 years of age
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

XDEMVY (lotilaner)

Products Affected

• XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including documentation of presence of mites upon examination of eyelashes by light microscopy or presence of collarettes on slit lamp examination, documentation of at least mild erythema of lid margin.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to ophthalmology and optometry
Coverage Duration	6 weeks
Other Criteria	The safety and efficacy of retreating with additional courses has not been fully described. For this reason, only one 6-week treatment course will be allowed every 365 days.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

XOSPATA (gilteritinib)

Products Affected

XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	Uncorrected hypokalemia or hypomagnesemia, coadministration with dual strong CYP3A/P-glycoprotein inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of FLT3 mutation, baseline serum potassium and magnesium levels, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

XURIDEN (uridine triacetate)

Products Affected

• XURIDEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of baseline CBC including neutrophil count and mean corpuscular volume, baseline urine orotic acid level.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, documentation of improvements or stabilization of urine orotic acid level, neutrophil count, and mean corpuscular volume is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

ZYKADIA (ceritinib)

Products Affected

• ZYKADIA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

INDEX

ACTIMMUNE 1	butalbital-apap-caffeine oral tablet 50-	
adapalene external cream236	325-40 mg	26
adapalene external gel 0.3 %236	butalbital-asa-caff-codeine	26
ADEMPAS ORAL TABLET 0.5 MG, 1 MG, 1.5	butalbital-aspirin-caffeine oral capsule	26
MG, 2 MG, 2.5 MG2	BYLVAY	27
AIMOVIG SUBCUTANEOUS SOLUTION	CABLIVI	29
AUTO-INJECTOR 140 MG/ML, 70 MG/ML38	CABOMETYX	30
AKEEGA 178	CALQUENCE ORAL TABLET	31
ALECENSA4	CAMZYOS	32
ALUNBRIG5	CAPLYTA ORAL CAPSULE 10.5 MG, 21 MG,	
alyq179	42 MG	.201
ambrisentan oral tablet 10 mg, 5 mg 179	CAPRELSA	34
apomorphine hcl subcutaneous 163	carglumic acid oral tablet soluble	35
AQNEURSA6	CERDELGA	
ARIKAYCE7	CHENODAL	40
ASCOMP-CODEINE26	CHOLBAM	41
AUGTYRO8	COBENFY	201
AUSTEDO ORAL TABLET 12 MG, 6 MG, 9	COBENFY STARTER PACK	201
MG9	COMETRIQ (100 MG DAILY DOSE) ORAL	
AUSTEDO XR ORAL TABLET EXTENDED	KIT 80 & 20 MG	42
RELEASE 24 HOUR 12 MG, 18 MG, 24 MG,	COMETRIQ (140 MG DAILY DOSE) ORAL	
30 MG, 36 MG, 42 MG, 48 MG, 6 MG9	KIT 3 X 20 MG & 80 MG	42
AUSTEDO XR PATIENT TITRATION ORAL	COMETRIQ (60 MG DAILY DOSE)	42
TABLET EXTENDED RELEASE THERAPY	COPIKTRA ORAL CAPSULE 15 MG, 25 MG	
PACK 12 & 18 & 24 & 30 MG9	CORLANOR ORAL SOLUTION	174
AUVELITY10	CORTROPHIN	44
AVMAPKI FAKZYNJA CO-PACK11	CORTROPHIN GEL	44
AYVAKIT ORAL TABLET 100 MG, 200 MG,	COTELLIC	46
25 MG, 300 MG, 50 MG12	CRESEMBA ORAL CAPSULE 186 MG	47
BALVERSA13	CYSTADROPS	48
BENLYSTA SUBCUTANEOUS16	CYSTARAN	48
BESREMI17	dalfampridine er	49
bexarotene external19	DANZITEN	144
bosentan oral tablet 125 mg, 62.5 mg179	dasatinib oral tablet 100 mg, 140 mg, 20	
BOSULIF ORAL CAPSULE 100 MG, 50 MG 22	mg, 50 mg, 70 mg, 80 mg	50
BOSULIF ORAL TABLET22	DAURISMO ORAL TABLET 100 MG, 25 MG	51
BRAFTOVI ORAL CAPSULE 75 MG24	deferasirox oral tablet soluble	52
BRUKINSA25	deferiprone	53
butalbital-acetaminophen oral tablet 50-	DIACOMIT	54
<i>325 mg</i> 26	dichlorphenamide	55
butalbital-apap-caff-cod26	diclofenac epolamine external	56
butalbital-apap-caffeine oral capsule 26	diclofenac sodium external gel 3 %	174
	digoxin oral tablet 250 mcg	57

dronabinol60 IBTROZI	
ui ui ui uui ii uu	91
DUPIXENT SUBCUTANEOUS SOLUTION icatibant acetate subcutaneous solution	7
AUTO-INJECTOR61 prefilled syringe	87
DUPIXENT SUBCUTANEOUS SOLUTION ICLUSIG	92
PREFILLED SYRINGE 200 MG/1.14ML, 300 IDHIFA	93
MG/2ML61 IMBRUVICA ORAL CAPSULE	94
eltrombopag olamine oral packet63 IMBRUVICA ORAL SUSPENSION	94
eltrombopag olamine oral tablet 12.5 mg, IMBRUVICA ORAL TABLET 140 MG, 280	
25 mg, 50 mg, 75 mg 63 MG, 420 MG	
EMGALITY38 IMKELDI	95
EMGALITY (300 MG DOSE)38 INBRIJA	163
ENSACOVE65 INCRELEX	135
EOHILIA66 INLYTA	98
EPIDIOLEX67 INQOVI	99
ERIVEDGE	100
ERLEADA178 INVEGA HAFYERA	101
eszopiclone	
everolimus oral tablet 10 mg, 2.5 mg, 5 SUSPENSION PREFILLED SYRINGE 273	
<i>mg, 7.5 mg</i> 69 MG/0.88ML, 410 MG/1.32ML, 546	
everolimus oral tablet soluble69 MG/1.75ML, 819 MG/2.63ML	101
EVRYSDI70 ITOVEBI ORAL TABLET 3 MG, 9 MG	103
ezetimibe-simvastatin oral tablet 10-80 mg ivabradine hcl oral tablet 5 mg, 7.5 mg.	174
205 IWILFIN	105
FABHALTA71 JAKAFI	106
FANAPT ORAL TABLET 1 MG, 10 MG, 12 javygtorjavygtor	199
MG, 2 MG, 4 MG, 6 MG, 8 MG201 JAYPIRCA	107
FANAPT TITRATION PACK A201 JOENJA	108
FANAPT TITRATION PACK B ORAL TABLET. 201 JUXTAPID ORAL CAPSULE 10 MG, 20 MG	G,
FANAPT TITRATION PACK C ORAL TABLET. 201 30 MG, 5 MG	109
FILSPARI73 KALYDECO73	37
FINTEPLA75 KERENDIA	111
FIRDAPSE76 ketoconazole oral	112
FOTIVDA77 KINERET SUBCUTANEOUS SOLUTION	
FRUZAQLA ORAL CAPSULE 1 MG, 5 MG78 PREFILLED SYRINGE	20
GAMMAGARD96 KISQALI (200 MG DOSE)	113
GAMMAGARD S/D LESS IGA96 KISQALI (400 MG DOSE)	113
GATTEX79 KISQALI (600 MG DOSE)	113
GAVRETO80 KOSELUGO	114
gefitinib81 KRAZATI	115
GILOTRIF82 lapatinib ditosylate	116
GOMEKLI85 LAZCLUZE ORAL TABLET 240 MG, 80 MG	
HAEGARDA88 lenalidomide	
HERNEXEOS89 LENVIMA (10 MG DAILY DOSE)	120

LENVIMA (12 MG DAILY DOSE) 120	nitisinone146
LENVIMA (14 MG DAILY DOSE) 120	NORDITROPIN FLEXPRO SUBCUTANEOUS
LENVIMA (18 MG DAILY DOSE) 120	SOLUTION PEN-INJECTOR 86
LENVIMA (20 MG DAILY DOSE) 120	NUBEQA
LENVIMA (24 MG DAILY DOSE) 120	NUCALA SUBCUTANEOUS SOLUTION
LENVIMA (4 MG DAILY DOSE)	AUTO-INJECTOR147
LENVIMA (8 MG DAILY DOSE)	NUCALA SUBCUTANEOUS SOLUTION
LEUKINE INJECTION SOLUTION	PREFILLED SYRINGE 100 MG/ML147
RECONSTITUTED121	NUCALA SUBCUTANEOUS SOLUTION
l-glutamine oral packet122	RECONSTITUTED147
lidocaine external patch 5 % 123	NUEDEXTA149
lidocan	NUPLAZID ORAL CAPSULE150
LIVTENCITY 124	NUPLAZID ORAL TABLET 10 MG150
LODOCO125	NURTEC38
LONSURF 127	OCTAGAM INTRAVENOUS SOLUTION 1
LORBRENA ORAL TABLET 100 MG, 25 MG. 128	GM/20ML, 10 GM/100ML, 10 GM/200ML,
LUMAKRAS 129	2 GM/20ML, 2.5 GM/50ML, 20
LYBALVI130	GM/200ML, 5 GM/100ML, 5 GM/50ML 96
LYNPARZA ORAL TABLET 131	ODOMZO151
LYTGOBI (12 MG DAILY DOSE)133	OFEV
LYTGOBI (16 MG DAILY DOSE)133	OGSIVEO ORAL TABLET 100 MG, 150 MG,
LYTGOBI (20 MG DAILY DOSE)133	50 MG153
MAVYRET134	OJEMDA154
megestrol acetate oral suspension 40	OJJAARA ORAL TABLET 100 MG, 150 MG,
mg/ml, 625 mg/5ml136	200 MG155
MEKINIST137	ONUREG
MEKTOVI24	OPSUMIT179
memantine hcl-donepezil hcl138	ORENITRAM
mifepristone oral tablet 300 mg139	ORENITRAM MONTH 1179
miglustat140	ORENITRAM MONTH 2179
MODEYSO141	ORENITRAM MONTH 3179
MOUNJARO SUBCUTANEOUS SOLUTION	ORIAHNN157
AUTO-INJECTOR 10 MG/0.5ML, 12.5	ORILISSA ORAL TABLET 150 MG, 200 MG 158
MG/0.5ML, 15 MG/0.5ML, 2.5 MG/0.5ML,	ORKAMBI37
5 MG/0.5ML, 7.5 MG/0.5ML83	ormalvi55
MYTESI142	ORSERDU
NAMZARIC ORAL CAPSULE EXTENDED	OTEZLA ORAL TABLET 30 MG20
RELEASE 24 HOUR 7-10 MG 138	OTEZLA ORAL TABLET THERAPY PACK 10 &
NAYZILAM174	20 & 30 MG20
NERLYNX143	OZEMPIC (0.25 OR 0.5 MG/DOSE)
NEXLETOL14	SUBCUTANEOUS SOLUTION PEN-INJECTOR
NEXLIZET14	2 MG/3ML83
nilotinib hcl144	OZEMPIC (1 MG/DOSE) SUBCUTANEOUS
NINLARO145	SOLUTION PEN-INJECTOR 4 MG/3ML 83

OZEMPIC (2 MG/DOSE)83	REXULTI ORAL TABLET 0.25 MG, 0.5 MG,	1
PANRETIN162	MG, 2 MG, 3 MG, 4 MG	201
pazopanib hcl165	REZDIFFRA	191
PEGASYS SUBCUTANEOUS SOLUTION 180	REZLIDHIA	192
MCG/ML174	REZUROCK	193
PEGASYS SUBCUTANEOUS SOLUTION	RIVFLOZA	194
PREFILLED SYRINGE174	ROMVIMZA	195
PEMAZYRE ORAL TABLET 13.5 MG, 4.5	ROZLYTREK	196
MG, 9 MG167	RUBRACA	197
PHEBURANE 168	RYBELSUS (FORMULATION R2) ORAL	
PIQRAY (200 MG DAILY DOSE) 169	TABLET 1.5 MG, 4 MG, 9 MG	83
PIQRAY (250 MG DAILY DOSE)	RYBELSUS ORAL TABLET 14 MG, 3 MG, 7	
PIQRAY (300 MG DAILY DOSE) 169	MG	83
pirfenidone oral tablet 267 mg, 801 mg 170	RYDAPT	198
POMALYST171	sajazir subcutaneous solution prefilled	
PRETOMANID	syringe	87
PREVYMIS ORAL TABLET173	sapropterin dihydrochloride oral packet	199
PRIVIGEN96	sapropterin dihydrochloride oral tablet	199
PROLASTIN-C INTRAVENOUS SOLUTION175	SCEMBLIX ORAL TABLET 100 MG, 20 MG,	
promethazine hcl oral solution 6.25	40 MG	200
<i>mg/5ml</i> 177	SECUADO TRANSDERMAL PATCH 24 HOL	JR
promethazine hcl oral tablet177	3.8 MG/24HR, 5.7 MG/24HR, 7.6	
promethazine hcl rectal suppository 12.5	MG/24HR	201
<i>mg, 25 mg</i> 177	SIGNIFOR	204
PROMETHEGAN RECTAL SUPPOSITORY 25	sildenafil citrate oral suspension	
MG, 50 MG177	reconstituted	179
QINLOCK181	sildenafil citrate oral tablet 20 mg	179
QULIPTA38	simvastatin oral tablet 80 mg	205
RADICAVA ORS	SIRTURO	206
RADICAVA ORS STARTER KIT182	SIVEXTRO	207
RECORLEV183	SKYCLARYS	208
REPATHA	sodium oxybate	161
REPATHA PUSHTRONEX SYSTEM185	sofosbuvir-velpatasvir	209
REPATHA SURECLICK185	SOHONOS	210
RETACRIT INJECTION SOLUTION 10000	SOMAVERT	211
UNIT/ML, 2000 UNIT/ML, 20000 UNIT/ML,	sorafenib tosylate	213
3000 UNIT/ML, 4000 UNIT/ML, 40000	SOTYKTU	20
UNIT/ML187	STELARA SUBCUTANEOUS SOLUTION 45	
RETEVMO ORAL TABLET 120 MG, 160 MG,	MG/0.5ML	20
40 MG, 80 MG188	STELARA SUBCUTANEOUS SOLUTION	
REVCOVI	PREFILLED SYRINGE	20
REVUFORJ ORAL TABLET 110 MG, 160 MG,	STIVARGA	214
25 MG190	SUCRAID	215
	sunitinib malate	216

SUNOSI ORAL TABLET 150 MG, 75 MG 217	TRUQAP ORAL TABLET	237
SYMDEKO 37	TRUQAP ORAL TABLET THERAPY PACK	237
TABRECTA218	TUKYSA ORAL TABLET 150 MG, 50 MG	238
tadalafil (pah)179	TURALIO ORAL CAPSULE 125 MG	239
tadalafil oral tablet 2.5 mg, 5 mg219	tyenne subcutaneous	20
TAFINLAR137	UBRELVY	38
TAGRISSO 222	UDENYCA	166
TAKHZYRO SUBCUTANEOUS SOLUTION88	UDENYCA ONBODY	166
TAKHZYRO SUBCUTANEOUS SOLUTION	ustekinumab subcutaneous	20
PREFILLED SYRINGE 300 MG/2ML88	UZEDY	240
TALZENNA ORAL CAPSULE 0.1 MG, 0.25	VALCHLOR	241
MG, 0.35 MG, 0.5 MG, 0.75 MG, 1 MG223	VALTOCO 10 MG DOSE	174
tasimelteon224	VALTOCO 15 MG DOSE NASAL LIQUID	
TAVNEOS225	THERAPY PACK 2 X 7.5 MG/0.1ML	174
tazarotene external cream236	VALTOCO 20 MG DOSE NASAL LIQUID	
tazarotene external gel236	THERAPY PACK 2 X 10 MG/0.1ML	174
TAZVERIK227	VALTOCO 5 MG DOSE	174
TENCON ORAL TABLET 50-325 MG 26	VANFLYTA	242
TEPMETKO228	VANRAFIA	243
teriparatide subcutaneous solution pen-	VELSIPITY	20
injector 560 mcg/2.24ml229	VENCLEXTA ORAL TABLET 10 MG, 100 M	1G,
testosterone transdermal gel 10 mg/act	50 MG	245
(2%), 12.5 mg/act (1%), 20.25 mg/1.25gm	VENCLEXTA STARTING PACK	245
(1.62%), 20.25 mg/act (1.62%), 25	VEOZAH	246
mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%),	VERQUVO ORAL TABLET 10 MG, 2.5 MG	, 5
50 mg/5gm (1%)231	MG	248
testosterone transdermal solution231	VERZENIO	250
TIBSOVO232	VIBERZI	251
tolvaptan oral tablet 15 mg, 30 mg 233	VIJOICE ORAL PACKET	252
tolvaptan oral tablet 15 mgtolvaptan	VIJOICE ORAL TABLET THERAPY PACK 12	25
(hyponatremia), 30 mgtolvaptan	MG, 200 & 50 MG, 50 MG	252
(hyponatremia)235	VITRAKVI ORAL CAPSULE 100 MG, 25 M	
tolvaptan oral tablet therapy pack 15 mg,	VITRAKVI ORAL SOLUTION	253
30 & 15 mg, 45 & 15 mg, 60 & 30 mg, 90 &	VIVJOA	254
<i>30 mg</i> 233	VIZIMPRO ORAL TABLET 15 MG, 30 MG,	45
torpenz69	MG	255
tretinoin external cream236	VONJO	256
tretinoin external gel 0.01 %, 0.025 %236	VOQUEZNA ORAL TABLET 10 MG, 20 MG	G 257
tridacaine ii123	VORANIGO ORAL TABLET 10 MG, 40 MG	i259
TRIKAFTA ORAL TABLET THERAPY PACK	VOWST	260
100-50-75 & 150 MG, 50-25-37.5 & 75 MG.37	VOYDEYA ORAL TABLET	261
TRIKAFTA ORAL THERAPY PACK	VOYDEYA ORAL TABLET THERAPY PACK.	261
TRULICITY SUBCUTANEOUS SOLUTION	VYNDAMAX	220
AUTO-INJECTOR83	VYNDAQEL	220

WEGOVY SUBCUTANEOUS SOLUTION	
AUTO-INJECTOR 0.25 MG/0.5ML, 0.5	
MG/0.5ML, 1 MG/0.5ML, 1.7 MG/0.75ML	,
2.4 MG/0.75ML	. 262
WELIREG	.264
WINREVAIR SUBCUTANEOUS KIT 2 X 45	
MG, 2 X 60 MG, 45 MG, 60 MG	.265
XALKORI	.267
XDEMVY	.268
XERMELO	.269
XIFAXAN ORAL TABLET 550 MG	. 270
XOLAIR	.272
XOSPATA	274
XPOVIO (100 MG ONCE WEEKLY) ORAL	
TABLET THERAPY PACK 50 MG	.275
XPOVIO (40 MG ONCE WEEKLY) ORAL	
TABLET THERAPY PACK 10 MG, 40 MG	. 275
XPOVIO (40 MG TWICE WEEKLY) ORAL	
TABLET THERAPY PACK 40 MG	.275
XPOVIO (60 MG ONCE WEEKLY) ORAL	
TABLET THERAPY PACK 60 MG	.275
XPOVIO (60 MG TWICE WEEKLY)	275
XPOVIO (80 MG ONCE WEEKLY) ORAL	
TABLET THERAPY PACK 40 MG	.275
XPOVIO (80 MG TWICE WEEKLY)	275
XTANDI	. 178
XURIDEN	.276
yargesa	
zaleplon	. 203
ZEJULA ORAL TABLET	277
ZELBORAF	46
ZILBRYSQ	
zolpidem tartrate oral tablet	.203
ZORYVE EXTERNAL CREAM 0.3 %	. 280
ZTALMY	281
ZURZUVAE	. 282
ZYDELIG	. 283
ZYKADIA ORAL TABLET	. 284