

## **PRIOR AUTHORIZATION CRITERIA**

This list is current as of 10/1/2025 and pertains to the following formularies:

2026 Independent Health's Medicare Advantage Standard Part D Formulary

Independent Health requires you (or your physician) to get prior authorization for certain drugs. This means that you will need to get approval from us before you fill your prescriptions. If you do not get approval, we may not cover the drug. These drugs are listed with a "PA" in the Requirements/Limits column on the formularies. This document contains the Prior Authorization requirements that are associated with our Medicare Advantage Part D formularies.

If you have any questions, please contact our Medicare Member Services Department at 1-800-665-1502 or, for TTY users 711, October 1<sup>st</sup> – March 31<sup>st</sup>: Monday through Sunday from 8 a.m. to 8 p.m., April 1<sup>st</sup> – September 30<sup>th</sup>: Monday through Friday from 8 a.m. to 8 p.m.

The formulary may change at any time. You will receive notice when necessary.

# ACTIMMUNE (interferon gamma-1b)

---

## Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# ADEMPAS (riociguat)

## Products Affected

- ADEMPAS ORAL TABLET 0.5 MG, 1 MG, 1.5 MG, 2 MG, 2.5 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, severe hepatic impairment (Child-Pugh class C), creatinine clearance below 15 mL/min or on dialysis, concurrent use with nitrates or nitric oxide donors in any form, concurrent use with phosphodiesterase inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use confirmed by right heart catheterization, submission of patient's WHO Group classification, mean pulmonary arterial pressure greater than 20 mm Hg at rest, pulmonary arterial wedge pressure less than or equal to 15 mm Hg, creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), and pregnancy status for female patients of childbearing potential. For pulmonary arterial hypertension (PAH, WHO Group 1), documentation of pulmonary vascular resistance (PVR) greater than 2 Woods units, submission of current or previous therapies used to treat the condition (see Other Criteria). For chronic thromboembolic pulmonary hypertension (CTEPH, WHO Group 4), confirmation of PVR greater than 3 Woods units, evidence of chronic pulmonary embolism on computed tomography or ventilation/perfusion (V/Q) scan, and attestation patient has inoperable disease or has persistent or recurrent disease after CTEPH surgery (pulmonary thromboendarterectomy).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology and pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization for PAH (WHO Group 1), the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to both (1) sildenafil or tadalafil and (2) ambrisentan or bosentan.

10/1/2025

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# ALECENSA (alectinib)

---

## Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# ALUNBRIG (brigatinib)

---

## Products Affected

- ALUNBRIG

PA Criteria	Criteria Details
Exclusion Criteria	Uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, baseline blood pressure reading, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# AUGTYRO (repotrectinib)

## Products Affected

- AUGTYRO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A4 inhibitors or inducers or P-glycoprotein inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For non-small cell lung cancer, submission of test confirming tumor is ROS1-positive. For other solid tumors, submission of test confirming tumor has a neurotrophic tyrosine receptor kinase (NTRK) gene fusion, attestation patient has progressed following treatment or patient has no satisfactory alternative therapy.
<b>Age Restrictions</b>	12 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## AUVELITY (dextromethorphan/bupropion)

---

### Products Affected

- AUVELITY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Seizure disorder, current or prior diagnosis of bulimia or anorexia nervosa, severe hepatic impairment, severe renal impairment, administration of monoamine oxidase inhibitors within 14 days of initiation
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient has been screened for and does not have bipolar disorder, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to psychiatry
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to two generic on-formulary antidepressants (e.g., bupropion, SSRI, SNRI).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025



## AYVAKIT (avapritinib)

### Products Affected

- AYVAKIT ORAL TABLET 100 MG, 200 MG, 25 MG, 300 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers or strong CYP3A inhibitors. For advanced or indolent systemic mastocytosis, platelet count below $50 \times 10^9/L$ .
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For gastrointestinal stromal tumor (GIST), submission of test result confirming presence of PDGFRA exon 18 mutation. For advanced or indolent systemic mastocytosis, submission of platelet count.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to allergy, hematology, immunology, and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## BALVERSA (erdafitinib)

### Products Affected

- BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	PD-1/PD-L1 inhibitor-eligible patients who have not received this therapy, coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of susceptible FGFR3 genetic alterations, submission of current or previous therapies used to treat the condition (see Other Criteria), pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	This drug is not recommended for the treatment of patients who are eligible for and have not received prior PD-1 or PD-L1 inhibitor therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# BEMPEDOIC ACID

## Products Affected

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant pravastatin utilization with doses above 40 mg/day, concomitant simvastatin utilization with doses above 20 mg/day, history of tendon disorders or rupture
<b>Required Medical Information</b>	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must (1) currently be using a statin (unless contraindicated) plus ezetimibe, or (2) have tried and failed to have an adequate response to or had an intolerance to (a) at least two statins or (b) one statin and ezetimibe. At least one statin previously tried and failed must be a hydrophilic statin. For each annual reauthorization, documentation that the patient remains on previously-used lipid-lowering therapies since the previous approval, unless there is documentation of a new contraindication or intolerance requiring discontinuation of a therapy (or therapies) since the previous approval, is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

10/1/2025

PA Criteria	Criteria Details
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# BENLYSTA (belimumab)

## Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe active central nervous system lupus, patients using other biologic medications or intravenous cyclophosphamide
<b>Required Medical Information</b>	Diagnosis of covered use, submission of current therapies used to treat the condition (see Other Criteria). For systemic lupus erythematosus, submission of autoantibody-positive test result for anti-nuclear antibodies (ANA) and/or anti-double-stranded DNA (anti-dsDNA).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to immunology, nephrology, and rheumatology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must be using standard therapy, defined as at least one of the following: systemic corticosteroids (e.g., prednisone), antimalarials (e.g., hydroxychloroquine), or immunosuppressants (e.g., azathioprine, methotrexate, mycophenolate mofetil). For each annual reauthorization, confirmation patient is still using some form of standard therapy (as defined above), unless contraindicated, is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

## BESREMI (ropeginterferon alfa-2b-njft)

### Products Affected

- BESREMI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History or presence of severe psychiatric disorders (including severe depression or suicidal ideation), history of presence of active serious or untreated autoimmune disease, moderate or severe hepatic impairment (Child-Pugh class B or C), immunosuppressed transplant recipients, severe or unstable cardiovascular disease (e.g., uncontrolled hypertension, NYHA class 2-4 congestive heart failure, serious cardiac arrhythmia, significant coronary artery stenosis, unstable angina), stroke or myocardial infarction within previous 6 months, severe renal impairment (eGFR less than 30 mL/min)
<b>Required Medical Information</b>	Diagnosis of covered use, submission of eGFR, documentation patient has tried and failed to have an adequate response to or had an intolerance/contraindication to hydroxyurea (HU), pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	If patient is taking HU, initially 12 weeks, then 1 year. If patient is not taking HU, 1 year.
<b>Other Criteria</b>	For the first reauthorization in patients using HU at the start of therapy, attestation patient has tapered completely off HU by the end of week 12 is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

10/1/2025

PA Criteria	Criteria Details
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# BEXAROTENE GEL

---

## Products Affected

- *bexarotene external*

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



# BIOLOGIC RESPONSE MODIFIERS

## Products Affected

- KINERET SUBCUTANEOUS SOLUTION MG/0.5ML
- PREFILLED SYRINGE
- OTEZLA ORAL TABLET 30 MG
- OTEZLA ORAL TABLET THERAPY PACK 10 & 20 & 30 MG
- SOTYKTU
- STELARA SUBCUTANEOUS SOLUTION 45
- STELARA SUBCUTANEOUS SOLUTION
- PREFILLED SYRINGE
- *tyenne subcutaneous*
- *ustekinumab subcutaneous*
- VELSIPITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria). For all drugs managed by this policy except Otezla and Velsipity, submission of baseline latent tuberculosis screening test (Mantoux tuberculin skin test [a.k.a. PPD test] or interferon-gamma release assay [IGRA]).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For initial authorization of a drug managed by this policy, the patient must have tried and failed to have an adequate response to or had an intolerance to at least two preferred agents (an adalimumab biosimilar, Cosentyx, Enbrel, Rinvoq, Skyrizi, an ustekinumab biosimilar, and Xeljanz/Xeljanz XR) for the indication submitted, where possible. For all drugs managed by this policy except Otezla and Velsipity, if TB screening test returns a positive result, coverage will be delayed until latent TB is treated. For each annual reauthorization, yearly TB screening test or chest X-ray required for patients who live in, work in, or travel to areas where TB exposure is likely while on treatment or for those who have previously had a positive TB screening test.
Indications	All FDA-approved Indications.
Off Label Uses	

10/1/2025

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

## BOSULIF (bosutinib)

### Products Affected

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inhibitors or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For accelerated or blast phase Ph+ CML, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	For initial authorization for accelerated or blast phase Ph+ CML, the patient must have either (1) tried and had an intolerance to dasatinib, imatinib, or nilotinib or (2) resistance to imatinib, defined as (a) failure to achieve or maintain any hematologic improvement within 4 weeks while on imatinib, or (b) failure to achieve a complete hematologic response by 3 months, cytogenetic response by 6 months or major cytogenetic response by 12 months, or (c) progression of disease after a previous cytogenetic or hematologic response, or (d) presence of a genetic mutation in the BCR-ABL gene associated with imatinib resistance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

10/1/2025

PA Criteria	Criteria Details
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

## BRAFTOVI/MEKTOVI (encorafenib/binimetinib)

---

### Products Affected

- BRAFTOVI ORAL CAPSULE 75 MG
- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of BRAF V600E (or V600K mutation for a melanoma diagnosis, pregnancy status for female patients of childbearing potential. For metastatic melanoma or metastatic non-small cell lung cancer, confirmation that encorafenib and binimetinib will be co-administered. For metastatic colorectal cancer, confirmation that encorafenib and cetuximab will be co-administered.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# BUTALBITAL-CONTAINING PRODUCTS IN OLDER PATIENTS

## Products Affected

- *butalbital-acetaminophen oral tablet 50-325 mg*
- *butalbital-apap-caff-cod*
- *butalbital-apap-caffeine oral capsule*
- *butalbital-apap-caffeine oral tablet 50-325-40 mg*
- *butalbital-asa-caff-codeine*
- *butalbital-aspirin-caffeine oral capsule*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation patient has tried and failed a preferred alternative such as ibuprofen or rizatriptan, or has contraindications to all alternatives.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

10/1/2025

## CABLIVI (caplacizumab-yhdp)

### Products Affected

- CABLIVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, confirmation drug will be given with plasma exchange and immunosuppressive therapy. If the coverage determination request is not for the patient's first use of caplacizumab, submission of previous aTTP recurrences while on caplacizumab.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology, hematology, and immunology
Coverage Duration	3 months
Other Criteria	If the coverage determination request is not for the patient's first use of caplacizumab, coverage will not be authorized if the patient has had more than 2 recurrences of aTTP while on therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# CALQUENCE (acalabrutinib)

---

## Products Affected

- CALQUENCE ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment, coadministration with strong CYP3A inhibitors
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



# CAMZYOS (mavacamten)

## Products Affected

- CAMZYOS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Left ventricular ejection fraction (LVEF) less than 55%, coadministration with strong CYP2C19 inhibitors, moderate or strong CYP2C19 inducers, moderate or strong CYP3A4 inducers, a non-dihydropyridine (DHP) calcium channel blocker (CCB) plus a beta-blocker, disopyramide, or ranolazine
<b>Required Medical Information</b>	Diagnosis of covered use including all three of the following: (1) attestation patient has exertional symptoms consistent with the definition of NYHA class II or III disease, (2) confirmation of left ventricular (LV) outflow tract obstruction gradient of at least 50 mm Hg either at rest, during Valsalva maneuver testing, or after exercise, and (3) confirmation of LV wall thickness of at least 15 mm or at least 13 mm if condition is familial, submission of current LVEF, current or previous therapies used to treat the condition (see Other Criteria), pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology
<b>Coverage Duration</b>	Initially 6 months, then 1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to both a beta-blocker and a non-DHP CCB. For each reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

10/1/2025

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# CAPRELSA (vandetanib)

## Products Affected

- CAPRELSA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of congenital long QT syndrome, torsades de pointes, uncompensated heart failure, or bradyarrhythmias, QTcF interval greater than 450 msec, hypocalcemia, hypokalemia, hypomagnesemia, coadministration with strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline serum potassium, calcium, magnesium, creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), ECG (or QT/QTcF interval), and pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

# CARGLUMIC ACID

---

## Products Affected

- *carglumic acid oral tablet soluble*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of elevated plasma ammonia level.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated plasma ammonia level since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# CERDELGA (eliglustat)

## Products Affected

- CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	Ultrarapid CYP2D6 metabolizers, pre-existing cardiac disease, moderate or severe hepatic impairment, long QT syndrome, coadministration with Class Ia or Class III antiarrhythmics. In patients who are poor or intermediate CYP2D6 metabolizers only, mild hepatic impairment.
Required Medical Information	Diagnosis of covered use, submission of CYP2D6 metabolizer status as detected by a test for determining CYP2D6 genotype, liver function testing or Child-Pugh score.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated liver function testing or Child-Pugh score since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# CFTR MODULATOR THERAPIES

## Products Affected

- KALYDECO 50-75 & 150 MG, 50-25-37.5 & 75 MG
- ORKAMBI • TRIKAFTA ORAL THERAPY PACK
- SYMDEKO
- TRIKAFTA ORAL TABLET THERAPY PACK 100-

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A inducers. For Trikafta, severe hepatic impairment.
<b>Required Medical Information</b>	Diagnosis of covered use, submission of cystic fibrosis (CF) mutation test confirming presence of CFTR gene mutations as indicated (see Other Criteria).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Initial authorization requires CF mutation test confirming presence of CFTR gene mutations as follows, by drug being requested: (a) for Kalydeco, a mutation predicted to be responsive to ivacaftor based on section 12.1 of the prescribing information, (b) for Orkambi, two copies of the F508del mutation, (c) for Symdeko, two copies of the F508del mutation or at least one mutation predicted to be responsive based on section 12.1 of the prescribing information, (d) for Trikafta, at least one mutation predicted to be responsive based on section 12.1 of the prescribing information or a responsive mutation based on in vitro data.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## CHENODAL (chenodiol)

---

### Products Affected

- CHENODAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, known hepatocyte dysfunction, bile duct abnormalities such as intrahepatic cholestasis, primary biliary cirrhosis, or sclerosing cholangitis, radiopaque stones, nonvisualizing gallbladder confirmed as nonvisualizing after 2 consecutive single doses of dye, compelling reasons for gallbladder surgery
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to gastroenterology and hepatology
<b>Coverage Duration</b>	24 months
<b>Other Criteria</b>	Safety beyond 24 months is not established and will not be authorized.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## CHOLBAM (cholic acid)

---

### Products Affected

- CHOLBAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of liver function testing.
Age Restrictions	
Prescriber Restrictions	Restricted to gastroenterology, hepatology, and pediatric gastroenterology
Coverage Duration	Initially 3 months, then 1 year
Other Criteria	For the first reauthorization, documentation of liver function improvement without complete biliary obstruction or persistent clinical or laboratory indications of worsening liver function or cholestasis is required. For each annual reauthorization, updated liver function testing since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



## COMETRIQ (cabozantinib)

### Products Affected

- COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment (Child-Pugh class C), uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of baseline blood pressure reading, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# COPIKTRA (duvelisib)

## Products Affected

- COPIKTRA ORAL CAPSULE 15 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Request as first- or second-line therapy, coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of at least two prior systemic therapies tried and failed, submission of pregnancy status for female patients of childbearing potential, attestation patient will receive prophylaxis for Pneumocystis jirovecii pneumonia (PJP) and, if necessary, cytomegalovirus.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# COTELLIC/ZELBORAF (cobimetinib/vemurafenib)

## Products Affected

- COTELLIC
- ZELBORAF

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For cobimetinib, coadministration with moderate or strong CYP3A inhibitors or inducers. For vemurafenib, electrolyte abnormalities that are not correctable, long QT syndrome, coadministration with drugs that prolong the QT interval.
<b>Required Medical Information</b>	Diagnosis of covered use including verification of BRAF V600 mutation as needed for diagnosis, submission of pregnancy status for female patients of childbearing potential. For patients using cobimetinib, submission of left ventricular ejection fraction (LVEF) with a requirement the baseline LVEF is greater than or equal to 50%. For patients using vemurafenib, submission of QTc interval with a requirement the QT interval is less than or equal to 500 msec.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## CRESEMBA (isavuconazonium)

---

### Products Affected

- CRESEMBA ORAL CAPSULE 186 MG

PA Criteria	Criteria Details
Exclusion Criteria	Familial short QT syndrome, coadministration with strong CYP3A inhibitors or inducers
Required Medical Information	Diagnosis of covered use.
Age Restrictions	6 years of age or older
Prescriber Restrictions	Restricted to hematology, infectious diseases, and oncology
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# CYSTEAMINE EYE DROPS

---

## Products Affected

- CYSTADROPS
- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, confirmation of corneal cysteine crystal deposits as seen on slit-lamp examination.
Age Restrictions	
Prescriber Restrictions	Restricted to metabolic diseases specialty, ophthalmology, and optometry
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# DALFAMPRIDINE

## Products Affected

- *dalfampridine er*

PA Criteria	Criteria Details
Exclusion Criteria	History of seizure, moderate or severe renal impairment (CrCl less than or equal to 50 mL/min)
Required Medical Information	Diagnosis of covered use, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), confirmation that patient is able to walk.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated creatinine clearance since the previous authorization and confirmation patient is able to walk is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# DASATINIB

## Products Affected

- *dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncorrected hypokalemia or hypomagnesemia, coadministration with proton pump inhibitors or H2 receptor antagonists
<b>Required Medical Information</b>	Diagnosis of covered use, submission of serum potassium and magnesium, pregnancy status for female patients of childbearing potential. For adults with resistance or intolerance to prior therapy, documentation of prior therapy.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

# DAURISMO (glasdegib)

---

## Products Affected

- DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, confirmation patient will also be receiving cytarabine as part of chemotherapeutic regimen. If patient is under 75 years of age, documentation of comorbidities that preclude use of intensive induction chemotherapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



# DEFERIPRONE

## Products Affected

- *deferiprone*

PA Criteria	Criteria Details
Exclusion Criteria	Absolute neutrophil count (ANC) below $1.5 \times 10^9/L$ , transfusional iron overload in myelodysplastic syndrome or Diamond Blackfan anemia
Required Medical Information	Diagnosis of covered use, submission of serum ferritin levels, ANC, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated ferritin level and ANC within last 3 months is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## DIACOMIT (stiripentol)

---

### Products Affected

- DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	Requests for monotherapy, moderate or severe renal impairment, moderate or severe hepatic impairment
Required Medical Information	Diagnosis of covered use, confirmation patient is also receiving clobazam.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# DICHLORPHENAMIDE

---

## Products Affected

- *dichlorphenamide*
- *ormalvi*

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of high dose aspirin, severe pulmonary disease limiting compensation to metabolic acidosis, hepatic insufficiency
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	Initially 2 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# DRONABINOL

## Products Affected

- *dronabinol*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use. If authorization is requested for treatment of nausea and vomiting associated with cancer therapy, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization for treatment of nausea and vomiting associated with cancer therapy, the patient must have tried and failed to have an adequate response to or had an intolerance to at least one 5-HT3 receptor antagonist (e.g., granisetron, ondansetron). If the medication is being administered related to cancer treatment and is a full replacement for intravenous administration of antiemetic therapy within 48 hours of cancer treatment, it is covered as a Part B benefit. To be eligible for Part B coverage, the prescribing physician must indicate this on the prescription.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

## ENSACOVE (ensartinib)

### Products Affected

- ENSACOVE

PA Criteria	Criteria Details
Exclusion Criteria	Prior ALK-inhibitor use, severe hepatic impairment, coadministration with moderate or strong CYP3A4 inhibitors or inducers or P-glycoprotein inhibitors
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, attestation patient has not previously received an ALK-inhibitor, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## EOHILIA (budesonide oral suspension)

### Products Affected

- EOHILIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation of upper endoscopy with biopsy showing at least 15 eosinophils per high-power field or 60 eosinophils/mm <sup>2</sup> , documentation of signs/symptoms, including but not limited to trouble swallowing, food sticking in esophagus, acid reflux, abdominal or chest pain, or nausea and vomiting, documentation patient has tried and failed at least an 8-week course of proton pump inhibitor therapy (i.e., patient has eosinophilic esophagitis unrelated to gastroesophageal reflux).
Age Restrictions	11 years of age or older
Prescriber Restrictions	Restricted to allergy, gastroenterology, immunology, and otolaryngology/otorhinolaryngology
Coverage Duration	12 weeks
Other Criteria	A maximum of one 12-week course will be allowed every 365 days.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

10/1/2025

## EPIDIOLEX (cannabidiol)

---

### Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## ERIVEDGE (vismodegib)

---

### Products Affected

- ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



# FILSPARI (sparsentan)

## Products Affected

- FILSPARI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, hepatic impairment, estimated glomerular filtration rate (eGFR) of at least 30 mL/min/1.73 m <sup>2</sup> , coadministration with renin-angiotensin system antagonists, endothelin receptor antagonists, proton pump inhibitors, or H <sub>2</sub> -receptor blockers
<b>Required Medical Information</b>	Diagnosis of primary IgA nephropathy confirmed by biopsy, submission of 24-hour urine protein of at least 1 g/day or 24-hour urine protein-to-creatinine ratio (UPCR) of at least 0.8 g/g, eGFR, liver function testing or Child-Pugh score, pregnancy status for female patients of childbearing potential, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to immunology and nephrology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must be currently using (or have a contraindication to the use of) a maximally-tolerated dose of an ACE inhibitor or ARB and tried and failed to have an adequate response to or had an intolerance to at least one immunosuppressant (e.g., azathioprine, mycophenolate, etc.). For each annual reauthorization, documentation of eGFR greater than or equal to 30 mL/min/1.73 m <sup>2</sup> and documentation of clinically relevant response to therapy, including either stabilization or improvement of UPCR or a reduction in 24-hour urine protein from baseline, is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

10/1/2025

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

## FINTEPLA (fenfluramine)

---

### Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	Administration of monoamine oxidase inhibitors within 14 days of initiation
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## GAVRETO (pralsetinib)

### Products Affected

- GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inhibitors, uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of RET gene fusion or mutation, baseline blood pressure reading, pregnancy status for female patients of childbearing potential. For thyroid cancer, attestation patient is radioactive iodine-refractory or ineligible.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# GEFITINIB

## Products Affected

- *gefinib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of EGFR exon 19 deletions or exon 21 (L858R) substitution mutations, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# GLP-1 AGONISTS

## Products Affected

- MOUNJARO SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.5ML, 12.5 MG/0.5ML, 15 MG/0.5ML, 2.5 MG/0.5ML, 5 MG/0.5ML, 7.5 MG/0.5ML
- OZEMPIC (0.25 OR 0.5 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 2 MG/3ML
- OZEMPIC (1 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 4 MG/3ML
- OZEMPIC (2 MG/DOSE)
- RYBELSUS (FORMULATION R2) ORAL TABLET 1.5 MG, 4 MG, 9 MG
- RYBELSUS ORAL TABLET 14 MG, 3 MG, 7 MG
- TRULICITY SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	Off-label use for weight management (see Other Criteria)
Required Medical Information	Diagnosis of type 2 diabetes confirmed through one of the following: (1) medical record, or (2) ICD-10 on medical claims, or (3) laboratory results (verifying a hemoglobin A1c greater than or equal to 6.5%, a fasting plasma glucose greater than or equal to 126 mg/dL, a 2-hour postprandial blood glucose greater than or equal to 200 mg/dL after an oral glucose tolerance test, or a random plasma blood glucose greater than or equal to 200 mg/dL combined with classic signs/symptoms of hyperglycemia or hyperglycemic crisis), attestation patient is not receiving another GLP-1 agonist for the treatment of any condition.
Age Restrictions	Age must be consistent with the prescribing information of the drug and condition being treated
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	These products will not be approved for weight management as this off-label use is currently excluded from coverage under Medicare Part D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

10/1/2025

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

10/1/2025

## GOMEKLI (mirdametinib)

---

### Products Affected

- GOMEKLI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



## HEREDITARY ANGIOEDEMA THERAPIES, ACUTE

---

### Products Affected

- *icabant acetate subcutaneous solution prefilled syringe*
- *sajazir subcutaneous solution prefilled syringe*

PA Criteria	Criteria Details
Exclusion Criteria	Requests for prophylactic hereditary angioedema therapy.
Required Medical Information	Diagnosis of covered use.
Age Restrictions	Age must be consistent with the prescribing information of the drug
Prescriber Restrictions	Restricted to allergy, dermatology, hematology, and immunology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# HEREDITARY ANGIOEDEMA THERAPIES, MAINTENANCE

## Products Affected

- HAEGARDA

PA Criteria	Criteria Details
Exclusion Criteria	Requests for acute hereditary angioedema (HAE) therapy (attacks)
Required Medical Information	Diagnosis of covered use, submission of objective or subjective documentation that prophylactic therapy is medically necessary, including, but not limited to activity of disease and disease burden, the frequency of HAE attacks, and quality of life.
Age Restrictions	Age must be consistent with the prescribing information of the drug
Prescriber Restrictions	Restricted to allergy, dermatology, hematology, and immunology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# HERNEXEOS (zongertinib)

## Products Affected

- HERNEXEOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including documentation the tumor is nonsquamous, submission of test confirming tumor is HER2-positive, pregnancy status for female patients of childbearing potential, documentation of at least one previous systemic therapy that was tried and failed.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# IBRANCE (palbociclib)

## Products Affected

- IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming primary tumor type is HR-positive, HER2-negative, attestation the treatment regimen will include concomitant use of an aromatase inhibitor or fulvestrant, pregnancy status for female patients of childbearing potential. For endocrine-resistant breast cancer, submission of genetic tumor testing confirming primary tumor type is PIK3CA-mutated, attestation patient will be taking palbociclib concurrently with inavolisib.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# IBTROZI (taletrectinib)

## Products Affected

- IBTROZI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A inhibitors or inducers, proton pump inhibitors, or H2 receptor antagonists
<b>Required Medical Information</b>	Diagnosis of covered use, submission of test confirming tumor is ROS1-positive, pregnancy status for female patients of childbearing potential, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to either crizotinib or entrectinib or have contraindications to both crizotinib and entrectinib.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# ICLUSIG (ponatinib)

## Products Affected

- ICLUSIG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Newly diagnosed chronic phase chronic myeloid leukemia (CP-CML), uncontrolled hypertension
<b>Required Medical Information</b>	Diagnosis of covered use, documentation of T315I mutation status (present or absent), submission of baseline blood pressure reading, pregnancy status for female patients of childbearing potential. For CP-CML that is not T315I-positive, documentation of resistance or intolerance to at least two prior kinase inhibitors. For Philadelphia chromosome-positive, T315I-negative acute lymphoblastic leukemia (monotherapy requests only), accelerated phase CML, or blast phase CML, attestation no other kinase inhibitors are indicated.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## IDHIFA (enasidenib)

---

### Products Affected

- IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of IDH2 mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## IMKELDI (imatinib)

### Products Affected

- IMKELDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of clinical rationale or documentation detailing why the patient cannot use imatinib oral tablets.
Age Restrictions	
Prescriber Restrictions	Restricted to allergy, dermatology, hematology, and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



## INQOVI (decitabine/cedazuridine)

---

### Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# INREBIC (fedratinib)

## Products Affected

- INREBIC

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, thiamine deficiency, coadministration with moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of thiamine level (see Other Criteria) and baseline platelet count, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to ruxolitinib. If baseline thiamine level is low, coverage will be delayed until thiamine is repleted.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# INVEGA INJECTABLE (paliperidone injectable suspension)

## Products Affected

- INVEGA HAFYERA MG/1.75ML, 819 MG/2.63ML
- INVEGA TRINZA INTRAMUSCULAR SUSPENSION PREFILLED SYRINGE 273 MG/0.88ML, 410 MG/1.32ML, 546

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis
<b>Required Medical Information</b>	Diagnosis of covered use. For the 3-month injection, documentation of at least 4 months' treatment with 1-month paliperidone palmitate extended-release injectable suspension. For the 6-month injection, documentation of at least 4 months' treatment with 1-month paliperidone palmitate extended-release injectable suspension or at least one 3-month injection of 3-month paliperidone palmitate extended-release injectable suspension.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

10/1/2025

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

10/1/2025

# ITOVEBI (inavolisib)

## Products Affected

- ITOVEBI ORAL TABLET 3 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming primary tumor type is HR-positive, HER2-negative, and PIK3CA-mutated, submission of current or previous therapies used to treat the condition (see Other Criteria), attestation that patient has locally advanced or metastatic disease, has not experienced disease progression on or following other PI3K inhibitors, and will be taking inavolisib concurrently with fulvestrant and palbociclib, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to endocrine therapy (e.g., tamoxifen or an aromatase inhibitor) and must not have received prior chemotherapy for metastatic breast cancer. In addition, documentation of clinical rationale why abemaciclib, palbociclib, or ribociclib combined with endocrine therapy is not suitable for the patient must be submitted.
Indications	All FDA-approved Indications.
Off Label Uses	

10/1/2025

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

## JAKAFI (ruxolitinib)

### Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	For myelofibrosis, platelet count less than $50 \times 10^9/L$
Required Medical Information	Diagnosis of covered use. For myelofibrosis, submission of baseline platelet count. For polycythemia vera, documented intolerance or inadequate response to hydroxyurea. For acute graft-versus-host disease, documented inadequate response to systemic corticosteroids. For chronic graft-versus-host-disease, documented failure of at least one previous line of systemic therapy.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology, oncology, and transplant specialty
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## JOENJA (leniolisib)

### Products Affected

- JOENJA

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, moderate or severe hepatic impairment (Child-Pugh class B or C)
Required Medical Information	Diagnosis of covered use including submission of test confirming presence of a pathogenic variant of either PIK3CD or PIK3R1, submission of liver function testing or Child-Pugh score, confirmation of negative pregnancy status for female patients of childbearing potential or attestation from physician patient is not pregnant and will be using a highly effective method of contraception, attestation patient is not currently using an immunosuppressive medication.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to specialists in genetic diseases or inborn errors of metabolism
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a clinical benefit (e.g., normalization of lymphocyte subsets, normalization of lymphadenopathy, reduction in spleen size, etc.) or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



# KETOCONAZOLE ORAL

## Products Affected

- *ketoconazole oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Acute or chronic liver disease, treatment of fungal meningitis or fungal infections of the skin or nails
<b>Required Medical Information</b>	Diagnosis of culture-proven systemic blastomycosis, coccidioidomycosis, histoplasmosis, chromomycosis, or paracoccidioidomycosis, submission of baseline ALT, AST, total bilirubin, alkaline phosphatase, prothrombin time and INR, prescriber attestation that the potential benefits of therapy outweigh the risks.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## KISQALI (ribociclib)

### Products Affected

- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Congenital long QT syndrome, QTcF interval greater than or equal to 450 msec at treatment initiation, uncorrected hypokalemia or hypomagnesemia, coadministration with strong CYP3A4 inducers or drugs that can prolong the QT interval
<b>Required Medical Information</b>	Diagnosis of covered use, submission of genetic tumor testing confirming the primary tumor type is HR-positive, HER2-negative, submission of QTcF interval, serum potassium and magnesium drawn within the previous 6 months, pregnancy status for female patients of childbearing potential, attestation that the treatment regimen will include concomitant use of an aromatase inhibitor or fulvestrant.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## KOSELUGO (selumetinib)

---

### Products Affected

- KOSELUGO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	Initiation: 2-17 years of age. Continuation: 2 years of age or older.
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	Selumetinib is indicated in pediatric patients and will not be approved for adults unless the patient started on the medication prior to 18 years of age.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# LAPATINIB

## Products Affected

- *lapatinib ditosylate*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncorrected hypokalemia or hypomagnesemia
<b>Required Medical Information</b>	Diagnosis of covered use, submission of genetic tumor testing confirming the primary tumor type is HER2-positive, submission of baseline potassium and magnesium levels, pregnancy status for female patients of childbearing potential, and attestation that the treatment regimen will include concomitant use of either capecitabine or letrozole. For patients who will be using lapatinib with capecitabine, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization for lapatinib in combination with capecitabine, the patient must have had prior therapy with at least an anthracycline, a taxane, and trastuzumab.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## LAZCLUZE (lazertinib)

---

### Products Affected

- LAZCLUZE ORAL TABLET 240 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations, attestation that the medication will be used in combination with amivantamab and will be given with anticoagulant prophylaxis for the first four months of therapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## LEUKINE (sargramostim, GM-CSF)

---

### Products Affected

- LEUKINE INJECTION SOLUTION  
RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# L-GLUTAMINE

---

## Products Affected

- *L-glutamine oral packet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of sickle cell disease, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	
Prescriber Restrictions	Restricted to hematology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must be using, tried and failed to have an adequate response to, or had an intolerance/contraindication to hydroxyurea.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

10/1/2025

# LIDOCAINE TRANSDERMAL PATCHES

---

## Products Affected

- *lidocaine external patch 5 %*
- *lidocan*
- *tridacaine ii*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



# LIVTENCITY (maribavir)

## Products Affected

- LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including a documented history of hematopoietic stem cell or solid organ transplant, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to hematology, infectious diseases, oncology, and transplant specialty
Coverage Duration	8 weeks
Other Criteria	For authorization, the patient must have tried and failed to have an adequate response to at least one of cidofovir, foscarnet, ganciclovir, or valganciclovir.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## LORBRENA (lorlatinib)

---

### Products Affected

- LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers, uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, baseline blood pressure, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## LYBALVI (olanzapine/samidorphan)

### Products Affected

- LYBALVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis, coadministration with opioids, levodopa, dopamine agonists, or strong CYP3A inducers, acute opioid withdrawal, end-stage renal disease
<b>Required Medical Information</b>	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology and psychiatry
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to both (1) generic olanzapine, including documentation showing a positive therapeutic benefit but unacceptable weight gain as a result of the drug, and (2) one other generic second-generation antipsychotic with low incidence of metabolic side effects (e.g., aripiprazole, lurasidone, ziprasidone).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# MAVYRET (glecaprevir/pibrentasvir)

---

## Products Affected

- MAVYRET

PA Criteria	Criteria Details
Exclusion Criteria	Moderate or severe hepatic impairment (Child-Pugh class B or C), coadministration with rifampin or atazanavir
Required Medical Information	Diagnosis of covered use, laboratory confirmation of hepatitis C virus (HCV) infection and HCV genotype.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Criteria for coverage duration will be applied consistent with current AASLD-IDSA guidance.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# MEGESTROL IN OLDER PATIENTS

---

## Products Affected

- *megestrol acetate oral suspension 40 mg/ml, 625 mg/5ml*

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	PA not required for hematology or oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## MEKINIST/TAFINLAR (trametinib/dabrafenib)

### Products Affected

- MEKINIST
- TAFINLAR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of BRAF V600E or V600K mutation, pregnancy status for female patients of childbearing potential. For anaplastic thyroid cancer, BRAF V600E-mutated solid tumors, low-grade glioma, and adjuvant BRAF V600E- and/or V600K-mutated melanoma indications, confirmation that trametinib and dabrafenib will be co-administered.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# MODEYSO (dordaviprone)

## Products Affected

- MODEYSO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming an H3K27M mutation, pregnancy status for female patients of childbearing potential, documentation of at least one previous therapy.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# NILOTINIB

## Products Affected

- DANZITEN
- *nilotinib hcl*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncorrected hypokalemia or hypomagnesemia, long QT syndrome, coadministration with drugs that prolong the QT interval, proton pump inhibitors, or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, confirmation of positive Philadelphia chromosome (Ph) status, baseline serum potassium and magnesium levels.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025



# NINLARO (ixazomib)

## Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, documentation that medication will be administered concomitantly with lenalidomide and dexamethasone, documentation of prior therapy regimen for multiple myeloma, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# NITISINONE

## Products Affected

- *nisinone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of succinylacetone in urine or plasma.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a clinical benefit, such as reductions in urine succinylacetone level, alpha-fetoprotein level, serum tyrosine level, or serum phenylalanine level, or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## NUPLAZID (pimavanserin)

### Products Affected

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Psychosis unrelated to Parkinson's disease psychosis, cardiac arrhythmias, symptomatic bradycardia, congenital QT prolongation, coadministration with moderate or strong CYP3A4 inducers or drugs that prolong the QT interval, uncorrected hypokalemia or hypomagnesemia
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline serum potassium and magnesium levels.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## ODOMZO (sonidegib)

---

### Products Affected

- ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, coadministration with strong CYP3A4 inhibitors or moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, attestation patient is not a candidate for surgery or radiation therapy or carcinoma has recurred following surgery or radiation therapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## OGSIVEO (nirogacestat)

---

### Products Affected

- OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inhibitors or inducers, proton pump inhibitors, or H2 receptor antagonists
Required Medical Information	Diagnosis of covered use with documentation of tumor progression, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology and sarcoma specialty
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## OJEMDA (tovorafenib)

### Products Affected

- OJEMDA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP2C8 inhibitors or inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of test confirming presence of BRAF V600 mutation or BRAF gene fusion or rearrangement, documentation of previous systemic therapy/therapies for pediatric low-grade glioma tried and failed with a minimum of one previous therapy necessary for approval, pregnancy status for female patients of childbearing potential. If genetic testing does not reveal a BRAF gene fusion or rearrangement, documentation of previous intolerance to, contraindication to, or other reason why the patient cannot use the combination of trametinib and dabrafenib.
<b>Age Restrictions</b>	Initiation: 21 years of age or younger (see Other Criteria)
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Tovorafenib is indicated as therapy in children and young adults and will not be approved for adults unless the patient started on the medication prior to 22 years of age.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

## OJJAARA (mometotinib)

---

### Products Affected

- OJJAARA ORAL TABLET 100 MG, 150 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	Active infection, uncontrolled acute or chronic liver disease
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## ONUREG (azacitidine)

### Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of myelodysplastic syndrome
Required Medical Information	Diagnosis of covered use, documentation patient achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy and attestation patient cannot complete intensive curative therapy, submission of absolute neutrophil count (with the requirement it is at least 500/mcL), submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



## ORIAHNN (elagolix/estradiol/norethindrone)

### Products Affected

- ORIAHNN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, known liver impairment or disease, known osteoporosis, undiagnosed abnormal uterine bleeding, women who are at increased risk of, have a history of, or currently have thrombotic or thromboembolic disorders (including women over 35 years of age who smoke and women with uncontrolled hypertension), current/history of breast cancer or other hormone-sensitive cancer
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient is premenopausal, submission of baseline blood pressure, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to endocrinology and gynecology
<b>Coverage Duration</b>	2 years (see Other Criteria)
<b>Other Criteria</b>	Use of this drug for more than 2 years increases risk of bone loss and requests for therapy for more than 2 years will not be approved.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

# OXYBATE SALT MEDICATIONS

## Products Affected

- *sodium oxybate*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with sedative hypnotics, diagnosis of insomnia
<b>Required Medical Information</b>	Diagnosis of covered use confirmed with documentation from a sleep study. For adults with excessive daytime sleepiness associated with narcolepsy, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	7 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology, psychiatry, and sleep medicine
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization for adults with excessive daytime sleepiness associated with narcolepsy, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to both (1) armodafinil or modafinil, and (2) solriamfetol. For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# PANRETIN (alitretinoin)

---

## Products Affected

- PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, instances where systemic Kaposi sarcoma therapy is required (more than 10 new Kaposi's sarcoma lesions in the prior month, symptomatic lymphedema, symptomatic pulmonary Kaposi sarcoma, or symptomatic visceral involvement)
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# PAZOPANIB

## Products Affected

- *pazopanib hcl*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, uncontrolled hypertension, uncorrected hypokalemia, hypocalcemia, or hypomagnesemia, coadministration with strong CYP3A4 inducers, proton pump inhibitors, H2-receptor antagonists, or drugs that can prolong the QT interval
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline blood pressure, serum potassium, calcium, and magnesium levels, pregnancy status for female patients of childbearing potential. For soft tissue sarcoma, submission of previous chemotherapy regimen(s).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

# PEGFILGRASTIM

## Products Affected

- UDENYCA
- UDENYCA ONBODY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of FDA-approved indication.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## PEMAZYRE (pemigatinib)

### Products Affected

- PEMAZYRE ORAL TABLET 13.5 MG, 4.5 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of either FGFR1 rearrangement or FGFR2 fusion or rearrangement depending on the indication, attestation patient has used previous systemic treatment for the indication, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# PRETOMANID

## Products Affected

- PRETOMANID

PA Criteria	Criteria Details
Exclusion Criteria	Inability to use bedaquiline or linezolid, drug-sensitive tuberculosis, coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, attestation pretomanid will be used in combination with bedaquiline and linezolid.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to infectious diseases and pulmonology
Coverage Duration	26 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# PREVYMIS (letermovir)

## Products Affected

- PREVYMIS ORAL TABLET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment (Child-Pugh class C), coadministration with ergot alkaloids, pimozide, or pitavastatin or simvastatin when coadministered with cyclosporine
<b>Required Medical Information</b>	Diagnosis of covered use, submission of day number post-transplant. For use after kidney transplant, documentation patient is high risk, defined as donor CMV seropositive/recipient CMV seronegative (D+/R-), submission of explanation why valganciclovir is contraindicated or cannot be used for prophylaxis.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology, infectious diseases, oncology, and transplant speciality
<b>Coverage Duration</b>	100 days post-HSCT or 200 days post-kidney transplant or post-HSCT at risk for late CMV infection
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025



# PROMETHAZINE IN OLDER PATIENTS

## Products Affected

- *promethazine hcl oral solution 6.25 mg/5ml*
- *promethazine hcl oral tablet*
- *promethazine hcl rectal suppository 12.5 mg, 25 mg*
- PROMETHEGAN RECTAL SUPPOSITORY 25 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. For allergic conditions, documentation must be submitted showing patient has tried and failed or had an inadequate response to a second-generation antihistamine.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	Promethazine is a potent anticholinergic considered high-risk in older patients due to risks of confusion, dry mouth, constipation, and decreased clearance with advanced age.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

10/1/2025

# PROSTATE CANCER ORAL MEDICATIONS

## Products Affected

- AKEEGA
- ERLEADA
- NUBEQA
- XTANDI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For Akeega, severe hepatic impairment (Child-Pugh class C), uncontrolled hypertension, uncontrolled hypokalemia
<b>Required Medical Information</b>	Diagnosis of covered use. For Nubeqa, submission of current or previous therapies used to treat the condition (see Other Criteria). For Akeega, submission of test confirming presence of deleterious BRCA mutation, Child-Pugh score or liver function testing, baseline blood pressure reading, and serum potassium level.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology and urology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization of Nubeqa, the patient must have tried and failed to have an adequate response or had an intolerance to both Erleada and Xtandi.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# PULMONARY HYPERTENSION MEDICATIONS

## Products Affected

- *alyq*
- *ambrisentan oral tablet 10 mg, 5 mg*
- *bosentan oral tablet 125 mg, 62.5 mg*
- OPSUMIT
- ORENITRAM
- ORENITRAM MONTH 1
- ORENITRAM MONTH 2
- ORENITRAM MONTH 3
- *sildenafil citrate oral suspension reconstituted*
- *sildenafil citrate oral tablet 20 mg*
- *tadalafil (pah)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For ambrisentan, bosentan, or Opsumit, pregnancy. For ambrisentan or Orenitram, moderate or severe hepatic impairment. For tadalafil, severe hepatic impairment or creatinine clearance below 30 mL/min or on hemodialysis. For ambrisentan, idiopathic pulmonary fibrosis.
<b>Required Medical Information</b>	Diagnosis of covered use confirmed by right heart catheterization, submission of mean pulmonary arterial pressure greater than 20 mm Hg at rest, pulmonary arterial wedge pressure less than or equal to 15 mm Hg, pulmonary vascular resistance greater than 2 Woods units, and attestation patient is WHO Group 1. For ambrisentan, bosentan, or Opsumit, submission of pregnancy status for female patients of childbearing potential. For Opsumit only, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	For all drugs in this policy except bosentan, 18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology and pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization of Opsumit, the patient must have tried and failed to have an adequate response to or had an intolerance to ambrisentan or bosentan.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

10/1/2025

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes

10/1/2025

## RETEVMO (selpercatinib)

### Products Affected

- RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	Uncontrolled hypertension, coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of RET gene fusion or mutation, baseline blood pressure reading, pregnancy status for female patients of childbearing potential. For RET fusion-positive thyroid cancer, attestation patient is radioactive iodine-refractory or ineligible. For solid tumors with a RET gene fusion, documentation of previous systemic therapy tried or reason why patient has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## REVCovi (elapegademase-lvlr)

### Products Affected

- REVCovi

PA Criteria	Criteria Details
Exclusion Criteria	Severe thrombocytopenia
Required Medical Information	Diagnosis of covered use confirmed by either biochemical testing showing less than 1% of adenosine deaminase (ADA) catalytic activity in red blood cells or genetic testing showing biallelic ADA pathogenic variants, submission of baseline plasma ADA level (if genetic testing was submitted as confirmation of diagnosis) and platelet count.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology, immunology, and specialists in genetic diseases
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, submission of updated plasma ADA level demonstrating an improvement from baseline and platelet count is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## REVUFORJ (revumenib)

### Products Affected

- REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	QTcF interval greater than 450 msec at treatment initiation, white blood cell (WBC) count greater than $25 \times 10^9/L$ , uncorrected hypokalemia or hypomagnesemia, coadministration with moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of test confirming presence of a lysine methyltransferase 2A gene (KMT2A) translocation that is not a 11q23 partial tandem duplication, submission of baseline QTcF interval, serum potassium, serum magnesium, baseline WBC count, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## REZLIDHIA (olutasidenib)

---

### Products Affected

- REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of IDH1 mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



## ROMVIMZA (vimseltinib)

---

### Products Affected

- ROMVIMZA

PA Criteria	Criteria Details
Exclusion Criteria	Active liver or biliary tract disease (including increased ALP), pre-existing increased serum transaminases, total or direct bilirubin greater than the upper limit of normal
Required Medical Information	Diagnosis of covered use (and documentation surgical intervention is not possible or practical), submission of serum transaminases, total and direct bilirubin, and ALP, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## ROZLYTREK (entrectinib)

### Products Affected

- ROZLYTREK

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers or drugs that prolong the QTc interval
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For non-small cell lung cancer, submission of test confirming presence of ROS1-positive tumor. For solid tumors, submission of evidence of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation and attestation tumor is metastatic or surgical resection/other systemic therapies are unsatisfactory treatment options.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# RYDAPT (midostaurin)

## Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For acute myeloid leukemia, submission of test confirming presence of FLT3 mutation, attestation patient will be receiving cytarabine and daunorubicin induction and cytarabine consolidation with midostaurin.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to allergy, hematology, and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# SAPROPTERIN

## Products Affected

- *javygtor*
- *sapropterin dihydrochloride oral packet*
- *sapropterin dihydrochloride oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of blood phenylalanine concentration.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For the first reauthorization, documentation of reduction in blood phenylalanine concentration from pre-treatment baseline is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## SECOND-GENERATION ANTIPSYCHOTICS

### Products Affected

- CAPLYTA ORAL CAPSULE 10.5 MG, 21 MG, 42 MG
- COBENFY
- COBENFY STARTER PACK
- FANAPT ORAL TABLET 1 MG, 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG
- FANAPT TITRATION PACK A
- FANAPT TITRATION PACK B ORAL TABLET
- FANAPT TITRATION PACK C ORAL TABLET
- REXULTI ORAL TABLET 0.25 MG, 0.5 MG, 1 MG, 2 MG, 3 MG, 4 MG
- SECUADO TRANSDERMAL PATCH 24 HOUR 3.8 MG/24HR, 5.7 MG/24HR, 7.6 MG/24HR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis. For Cobenfy only, moderate to severe renal impairment, urinary retention, gastric retention, hepatic impairment, untreated narrow-angle glaucoma.
<b>Required Medical Information</b>	Diagnosis of covered use. For schizophrenia, an indication related to bipolar disorder type I, or for Rexulti for major depressive disorder, submission of current or previous therapies used to treat the condition (see Other Criteria). For Cobenfy only, submission of estimated glomerular filtration rate with the requirement it is at least 60 mL/min, attestation patient does not have hepatic impairment.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to psychiatry
<b>Coverage Duration</b>	1 year

10/1/2025

PA Criteria	Criteria Details
<b>Other Criteria</b>	For initial authorization for schizophrenia or acute treatment of manic/mixed episodes of bipolar I disorder, the patient must have tried and failed to have an adequate response to or had an intolerance to aripiprazole and at least one other generic second-generation atypical antipsychotic. For initial authorization of Caplyta for depressive episodes associated with bipolar I disorder, the patient must have tried and failed to have an adequate response to or had an intolerance to at least two of the following drugs: cariprazine, lurasidone, olanzapine, or quetiapine. For Rexulti as an adjunctive therapy to antidepressants for major depressive disorder, the patient must have tried and failed to have an adequate response to or had an intolerance to aripiprazole and cariprazine or quetiapine.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

## SEDATIVE HYPNOTICS

### Products Affected

- *eszopiclone*
- *zaleplon*
- *zolpidem tartrate oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use, documentation at least two of the following medications were tried and deemed ineffective or intolerable: Belsomra, ramelteon, and trazodone.
<b>Age Restrictions</b>	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Sedative hypnotic medications are high-risk medications in older patients due to increased risks of cognitive impairment, delirium, unsteady gait, syncope, falls, fractures, and motor vehicle accidents.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# SOFOSBUVIR/VELPATASVIR

## Products Affected

- *sofosbuvir-velpatasvir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, laboratory confirmation of hepatitis C virus (HCV) infection and HCV genotype, attestation that patients with decompensated cirrhosis will receive concomitant ribavirin therapy unless ribavirin therapy is contraindicated.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



# SOMAVERT (pegvisomant)

## Products Affected

- SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including attestation that surgery or radiation was not curative or is not an option, submission of baseline IGF-1, submission of baseline liver function testing (LFT) including bilirubin with the requirement liver transaminases are less than or equal to 3 times the upper limit of normal (ULN). If liver transaminases are greater than 3 times ULN, submission of the cause of liver dysfunction determined through a comprehensive workup.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to endocrinology
Coverage Duration	1 year
Other Criteria	For the first reauthorization, submission of updated IGF-1 level demonstrating an improvement from baseline, LFT showing liver transaminases below 5 times the ULN, and attestation patient does not have signs or symptoms of liver injury (e.g., jaundice, elevated bilirubin level or bilirubinuria, fatigue, nausea, vomiting, right upper quadrant pain, ascites, unexplained edema, easy bruisability) is required. For each annual reauthorization, documented improvement/maintenance of IGF-1 level is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

10/1/2025

PA Criteria	Criteria Details
Prerequisite Therapy Required	No

10/1/2025

# SORAFENIB

## Products Affected

- *sorafenib tosylate*

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome, coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For differentiated thyroid cancer, attestation patient is radioactive iodine-refractory or ineligible.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# TABRECTA (capmatinib)

---

## Products Affected

- TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of MET exon 14 skipping mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# TAGRISSO (osimertinib)

---

## Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of EGFR exon 19 deletions, exon 21 L858R mutations, or T790M mutations, pregnancy status for female patients of childbearing potential. For EGFR T790M mutation-positive NSCLC, submission of previous EGFR tyrosine kinase inhibitor therapy used for indication.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# TALZENNA (talazoparib)

## Products Affected

- TALZENNA ORAL CAPSULE 0.1 MG, 0.25 MG, 0.35 MG, 0.5 MG, 0.75 MG, 1 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For breast cancer, submission of test results confirming germline BRCA mutation-positive, human epidermal growth factor receptor 2 (HER2) negative disease. For prostate cancer, submission of test results confirming HRR gene-mutated disease, confirmation talazoparib will be used in combination with enzalutamide.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# TASIMELTEON

## Products Affected

- *tasimelteon*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, coadministration with strong CYP1A2 inhibitors or CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use. For non-24-hour sleep-wake disorder, attestation patient is totally blind. For Smith-Magenis Syndrome, documentation of genetic testing results confirming chromosome 17p11.2 deletion.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to neurology and sleep medicine
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## TEPMETKO (tepotinib)

### Products Affected

- TEPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of MET exon 14 skipping mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



# TERIPARATIDE

## Products Affected

- *teriparade subcutaneous solution pen-injector 560 mcg/2.24ml*

PA Criteria	Criteria Details
Exclusion Criteria	Pre-existing hypercalcemia, underlying hypercalcemic disorder (such as primary hyperparathyroidism), patients with an increased risk of osteosarcoma (such as those with Paget's disease)
Required Medical Information	Diagnosis of covered use where "high risk for fracture" is defined as documentation of (1) a history of fracture of the hip or vertebra regardless of bone mineral density (BMD), or (2) a history of fracture of the proximal humerus, pelvis, or distal forearm and T-score between -1.0 and -2.5, or (3) T-score less than or equal to -2.5 at the total hip, femoral neck, spine, or distal third of the radius, or (4) T-score between -1.0 and -2.5 at the total hip, femoral neck, spine, or distal third of the radius, and (a) a 10-year probability of hip fracture as assessed by FRAX score of at least 3%, or (b) a 10-year probability of a major osteoporosis-related fracture as assessed by FRAX score of at least 20%, submission of baseline serum calcium, postmenopausal status, current or previous therapies used to treat the condition (see Other Criteria), number of total months of all prior use of parathyroid hormone analogs and parathyroid hormone related peptides.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	2 years unless patient is at high risk for fracture after 2 years of therapy (see Other Criteria)

10/1/2025

PA Criteria	Criteria Details
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to at least one bisphosphonate. Therapeutic failure is defined as either a fracture or a decrease in BMD while using a bisphosphonate for at least 3 months. Use of parathyroid hormone analogs and/or parathyroid hormone related peptides for more than 2 years during a patient's lifetime is generally not recommended. For annual reauthorization beyond 2 years, submission of updated serum calcium since initial authorization and evidence the patient remains at high risk for fracture as defined in the Required Medical Information section is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

# TOLVAPTAN (HYPONATREMIA)

## Products Affected

- *tolvaptan oral tablet 15 mg* *tolvaptan (hyponatremia)*, *30 mg* *tolvaptan (hyponatremia)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Underlying liver disease, need to raise serum sodium acutely, inability to sense or respond to thirst, hypovolemia, anuria, coadministration with strong CYP3A inhibitors or inducers or desmopressin
<b>Required Medical Information</b>	Diagnosis of covered use, submission of evidence of clinically significant hyponatremia, defined as (1) serum sodium less than 125 mEq/L, or (2) serum sodium less than 135 mEq/L that is symptomatic and has resisted correction with fluid restriction.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	30 days (see Other Criteria)
<b>Other Criteria</b>	Treatment should be initiated in a setting where serum sodium can be monitored closely. Treatment is limited to 30 days to prevent liver injury. This formulation of tolvaptan will not be approved for autosomal dominant polycystic kidney disease (ADPKD) as it is not indicated for ADPKD and the tolvaptan formulation approved for ADPKD has a mandatory REMS program, making it only available through a restricted distribution program.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

# TRUQAP (capivasertib)

## Products Affected

- TRUQAP ORAL TABLET
- TRUQAP ORAL TABLET THERAPY PACK

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming that the primary tumor type is HR-positive, HER2-negative and has at least one PIK3CA, AKT1, and/or PTEN mutation, submission of pregnancy status for female patients of childbearing potential, attestation patient has (1) progressed on at least one endocrine-based regimen in the metastatic setting, or (2) has recurrent disease on or within 12 months of completing adjuvant therapy, attestation drug will be given with fulvestrant.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# TURALIO (pexidartinib)

## Products Affected

- TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	Active liver or biliary tract disease (including increased ALP), pre-existing increased serum transaminases, total or direct bilirubin greater than the upper limit of normal, coadministration with other hepatotoxic medications, strong CYP3A inducers, or proton pump inhibitors
Required Medical Information	Diagnosis of covered use (and documentation surgical intervention is not possible or practical), documentation of patient's severe morbidity or functional limitations, submission of serum transaminases, total and direct bilirubin, and ALP, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## VALCHLOR (mechlorethamine)

---

### Products Affected

- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	Use as initial therapy
Required Medical Information	Diagnosis of covered use, submission of previous skin-directed therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# VANFLYTA (quizartinib)

## Products Affected

- VANFLYTA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Requests for maintenance monotherapy after allogeneic hematopoietic stem cell transplant, uncorrected hypokalemia or hypomagnesemia, long QT syndrome, QTcF interval greater than 450 msec at treatment initiation, coadministration with moderate or strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use including submission of test confirming presence of FLT3 internal tandem duplication-positive mutation, submission of QTcF interval, baseline serum potassium and magnesium levels, and pregnancy status for female patients of childbearing potential, attestation patient does not have history of ventricular arrhythmias or torsades de pointes.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

# VANRAFIA (atrasentan)

## Products Affected

- VANRAFIA

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, severe hepatic impairment, coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of primary IgA nephropathy confirmed by biopsy, submission of 24-hour urine protein of at least 1 g/day or 24-hour urine protein-to-creatinine ratio (UPCR) of at least 0.8 g/g, liver function testing or Child-Pugh score, pregnancy status for female patients of childbearing potential, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to immunology and nephrology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must be currently using (or have a contraindication to the use of) a maximally-tolerated dose of an ACE inhibitor or ARB and tried and failed to have an adequate response to or had an intolerance to at least one immunosuppressant (e.g., azathioprine, mycophenolate, etc.). For each annual reauthorization, documentation of clinically relevant response to therapy, including either stabilization or improvement of UPCR or a reduction in 24-hour urine protein from baseline, is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

10/1/2025



PA Criteria	Criteria Details
<b>Prerequisite Therapy Required</b>	Yes

10/1/2025

## VENCLEXTA (venetoclax)

---

### Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers. For CLL/SLL, coadministration with strong CYP3A inhibitors at treatment initiation and initial dosage titration.
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## VERZENIO (abemaciclib)

### Products Affected

- VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers or ketoconazole
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming that the primary tumor type is HR-positive, HER2-negative, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# VIJOICE (alpelisib)

## Products Affected

- VIJOICE ORAL PACKET
- VIJOICE ORAL TABLET THERAPY PACK 125 MG, 200 & 50 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use including at least one target lesion on imaging with requesting provider attestation patient has severe or life-threatening disease, submission of test confirming presence of mutation in PIK3CA gene, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to specialists in genetic diseases or inborn errors of metabolism
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a symptomatic or clinical benefit (e.g., reductions in target lesion size, pain, vascular malformations, limb enlargements, etc.), or maintenance of improvement previously achieved, is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# VITRAKVI (larotrectinib)

---

## Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of evidence of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation, attestation tumor is metastatic or surgical resection and other systemic therapies are unsatisfactory treatment options, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## VIZIMPRO (dacomitinib)

---

### Products Affected

- VIZIMPRO ORAL TABLET 15 MG, 30 MG, 45 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with a proton pump inhibitor
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of EGFR exon 19 deletions or exon 21 L858R substitution mutations, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## VONJO (pacritinib)

### Products Affected

- VONJO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Moderate or severe hepatic impairment (Child-Pugh class B or C), estimated glomerular filtration rate (eGFR) less than 30 mL/min, QTc interval greater than 480 msec at baseline, uncorrected hypokalemia, coadministration with strong CYP3A4 inducers or strong CYP3A4 inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of platelet count, serum potassium level, eGFR, and QTc interval, Child-Pugh score, documentation from a physical exam patient has splenomegaly.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## VORANIGO (vorasidenib)

### Products Affected

- VORANIGO ORAL TABLET 10 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of IDH1 or IDH2 mutation, pregnancy status for female patients of childbearing potential, attestation patient has had at least one prior surgery (biopsy, sub-total resection, or gross total resection).
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025



## VOWST (fecal microbiota spores, live-brpk)

### Products Affected

- VOWST

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use with the requirement patient is being treated after at least 2 recurrent (3 total) Clostridioides difficile infections (confirmation of pathogen with stool test or other confirmatory test), submission of time of last planned dose of antibiotic for latest recurrent C. difficile infection and attestation patient will be using a bowel cleanse the evening prior to starting Vowst, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 course (3 days)
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had a contraindication to fecal microbiota, live-jslm rectal suspension.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	Yes
<b>Prerequisite Therapy Required</b>	No

10/1/2025

## XALKORI (crizotinib)

### Products Affected

- XALKORI

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome, coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming tumor is ALK or ROS1-positive, pregnancy status for female patients of childbearing potential.
Age Restrictions	For ALK-positive systemic anaplastic large cell lymphoma only, 1 year of age to 21 years of age
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# XDEMVY (lotilaner)

## Products Affected

- XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including documentation of presence of mites upon examination of eyelashes by light microscopy or presence of collarettes on slit lamp examination, documentation of at least mild erythema of lid margin.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to ophthalmology and optometry
Coverage Duration	6 weeks
Other Criteria	The safety and efficacy of retreating with additional courses has not been fully described. For this reason, only one 6-week treatment course will be allowed every 365 days.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# XOSPATA (gilteritinib)

## Products Affected

- XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	Uncorrected hypokalemia or hypomagnesemia, coadministration with dual strong CYP3A/P-glycoprotein inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of FLT3 mutation, baseline serum potassium and magnesium levels, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

# XPOVIO (selinexor)

## Products Affected

- XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 50 MG
- XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 10 MG, 40 MG
- XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (80 MG TWICE WEEKLY)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For multiple myeloma in combination with bortezomib and dexamethasone, documented failure of at least one previous therapy. For relapsed or refractory multiple myeloma, documented failure of at least four previous lines of systemic therapy including at least two proteasome inhibitors, at least two immunomodulatory agents, and an anti-CD38 monoclonal antibody. For relapsed or refractory diffuse large B-cell lymphoma, documented failure of at least two previous lines of systemic therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## XURIDEN (uridine triacetate)

---

### Products Affected

- XURIDEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of baseline CBC including neutrophil count and mean corpuscular volume, baseline urine orotic acid level.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, documentation of improvements or stabilization of urine orotic acid level, neutrophil count, and mean corpuscular volume is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

## ZYKADIA (ceritinib)

---

### Products Affected

- ZYKADIA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

10/1/2025

10/1/2025



## INDEX

ACTIMMUNE .....	1	<i>butalbital-asa-caff-codeine</i> .....	27
<i>adapalene external cream</i> .....	214	<i>butalbital-aspirin-caffeine oral capsule</i> .....	27
<i>adapalene external gel 0.3 %</i> .....	214	CABLIVI .....	28
ADEMPAS ORAL TABLET 0.5 MG, 1 MG, 1.5 MG, 2 MG, 2.5 MG .....	2	CABOMETYX .....	29
AIMOVIG SUBCUTANEOUS SOLUTION		CALQUENCE ORAL TABLET .....	30
AUTO-INJECTOR 140 MG/ML, 70 MG/ML .....	37	CAMZYOS .....	31
AKEEGA .....	160	CAPLYTA ORAL CAPSULE 10.5 MG, 21 MG, 42 MG .....	181
ALECENSA .....	4	CAPRELSA .....	33
ALUNBRIG .....	5	<i>carglumic acid oral tablet soluble</i> .....	34
<i>alyq</i> .....	161	CERDELGA .....	35
<i>ambrisentan oral tablet 10 mg, 5 mg</i> .....	161	CHENODAL .....	39
<i>apomorphine hcl subcutaneous</i> .....	6	CHOLBAM .....	40
AQNEURSA .....	7	COBENFY .....	181
ARIKAYCE .....	8	COBENFY STARTER PACK .....	181
AUGTYRO .....	9	COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG .....	41
AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG .....	10	COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG .....	41
AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HOUR 12 MG, 18 MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 MG, 6 MG .....	10	COMETRIQ (60 MG DAILY DOSE) .....	41
AUSTEDO XR PATIENT TITRATION ORAL TABLET EXTENDED RELEASE THERAPY PACK 12 & 18 & 24 & 30 MG .....	10	COPIKTRA ORAL CAPSULE 15 MG, 25 MG ...	42
AUVELITY .....	11	COTELLIC .....	43
AVMAPKI FAKZYNJA CO-PACK .....	12	CRESEMBA ORAL CAPSULE 186 MG .....	44
AYVAKIT ORAL TABLET 100 MG, 200 MG, 25 MG, 300 MG, 50 MG .....	13	CYSTADROPS .....	45
BALVERSA .....	14	CYSTARAN .....	45
BENLYSTA SUBCUTANEOUS .....	17	<i>dalfampridine er</i> .....	46
BESREMI .....	18	DANZITEN .....	131
<i>bexarotene external</i> .....	20	<i>dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg</i> .....	47
<i>bosentan oral tablet 125 mg, 62.5 mg</i> .....	161	DAURISMO ORAL TABLET 100 MG, 25 MG ..	48
BOSULIF ORAL CAPSULE 100 MG, 50 MG ...	23	<i>deferiprone</i> .....	49
BOSULIF ORAL TABLET .....	23	DIACOMIT .....	50
BRAFTOVI ORAL CAPSULE 75 MG .....	25	<i>dichlorphenamide</i> .....	51
BRUKINSA .....	26	<i>diclofenac sodium external gel 3 %</i> .....	193
<i>butalbital-acetaminophen oral tablet 50-325 mg</i> .....	27	<i>digoxin oral tablet 250 mcg</i> .....	52
<i>butalbital-apap-caff-cod</i> .....	27	DOPTELET ORAL TABLET 20 MG .....	53
<i>butalbital-apap-caffeine oral capsule</i> .....	27	<i>dronabinol</i> .....	55
<i>butalbital-apap-caffeine oral tablet 50-325-40 mg</i> .....	27	DUPIXENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR .....	56
		DUPIXENT SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 200 MG/1.14ML, 300 MG/2ML .....	56
		<i>eltrombopag olamine oral packet</i> .....	58

<i>eltrombopag olamine oral tablet 12.5 mg, 25 mg, 50 mg, 75 mg</i> .....	58	IMKELDI.....	87
EMGALITY.....	37	INCRELEX.....	123
EMGALITY (300 MG DOSE).....	37	INLYTA.....	90
ENSACOVE.....	60	INQOVI.....	91
EOHILIA.....	61	INREBIC.....	92
EPIDIOLEX.....	62	INVEGA HAFYERA.....	93
ERIVEDGE.....	63	INVEGA TRINZA INTRAMUSCULAR SUSPENSION PREFILLED SYRINGE 273 MG/0.88ML, 410 MG/1.32ML, 546 MG/1.75ML, 819 MG/2.63ML.....	93
<i>eszopiclone</i> .....	183	ITOVEBI ORAL TABLET 3 MG, 9 MG.....	95
<i>everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg</i> .....	64	<i>ivabradine hcl oral tablet 5 mg, 7.5 mg</i> .....	156
<i>everolimus oral tablet soluble</i> .....	64	IWILFIN.....	97
EVRYSDI.....	65	JAKAFI.....	98
<i>ezetimibe-simvastatin oral tablet 10-80 mg</i> .....	185	<i>javygtor</i> .....	179
FANAPT ORAL TABLET 1 MG, 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG.....	181	JAYPIRCA.....	99
FANAPT TITRATION PACK A.....	181	JOENJA.....	100
FANAPT TITRATION PACK B ORAL TABLET.....	181	KALYDECO.....	36
FANAPT TITRATION PACK C ORAL TABLET.....	181	KERENDIA.....	101
FILSPARI.....	66	<i>ketoconazole oral</i> .....	102
FINTEPLA.....	68	KINERET SUBCUTANEOUS SOLUTION PREFILLED SYRINGE.....	21
FIRDAPSE.....	69	KISQALI (200 MG DOSE).....	103
FOTIVDA.....	70	KISQALI (400 MG DOSE).....	103
FRUZAQLA ORAL CAPSULE 1 MG, 5 MG.....	71	KISQALI (600 MG DOSE).....	103
GAMMAGARD.....	88	KOSELUGO.....	104
GAMMAGARD S/D LESS IGA.....	88	KRAZATI.....	105
GAVRETO.....	72	<i>lapatinib ditosylate</i> .....	106
<i>gefitinib</i> .....	73	LAZCLUZE ORAL TABLET 240 MG, 80 MG.....	107
GILOTRIF.....	74	<i>lenalidomide</i> .....	108
GOMEKLI.....	77	LENVIMA (10 MG DAILY DOSE).....	110
HAEGARDA.....	80	LENVIMA (12 MG DAILY DOSE).....	110
HERNEXEOS.....	81	LENVIMA (14 MG DAILY DOSE).....	110
IBRANCE.....	82	LENVIMA (18 MG DAILY DOSE).....	110
IBTROZI.....	83	LENVIMA (20 MG DAILY DOSE).....	110
<i>icatibant acetate subcutaneous solution</i> <i>prefilled syringe</i> .....	79	LENVIMA (24 MG DAILY DOSE).....	110
ICLUSIG.....	84	LENVIMA (4 MG DAILY DOSE).....	110
IDHIFA.....	85	LENVIMA (8 MG DAILY DOSE).....	110
IMBRUVICA ORAL CAPSULE.....	86	LEUKINE INJECTION SOLUTION RECONSTITUTED.....	111
IMBRUVICA ORAL SUSPENSION.....	86	<i>l-glutamine oral packet</i> .....	112
IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG.....	86	<i>lidocaine external patch 5 %</i> .....	113
		<i>lidocan</i> .....	113
		LIVTENCITY.....	114

LONSURF .....	115	OCTAGAM INTRAVENOUS SOLUTION 1	
LORBRENA ORAL TABLET 100 MG, 25 MG .....	116	GM/20ML, 10 GM/100ML, 10 GM/200ML,	
LUMAKRAS .....	117	2 GM/20ML, 2.5 GM/50ML, 20	
LYBALVI .....	118	GM/200ML, 5 GM/100ML, 5 GM/50ML .....	88
LYNPARZA ORAL TABLET .....	119	ODOMZO .....	138
LYTGOBI (12 MG DAILY DOSE) .....	121	OFEV .....	139
LYTGOBI (16 MG DAILY DOSE) .....	121	OGSIVEO ORAL TABLET 100 MG, 150 MG,	
LYTGOBI (20 MG DAILY DOSE) .....	121	50 MG .....	140
MAVYRET .....	122	OJEMDA .....	141
<i>megestrol acetate oral suspension 40</i>		OJJAARA ORAL TABLET 100 MG, 150 MG,	
<i>mg/ml, 625 mg/5ml .....</i>	124	200 MG .....	142
MEKINIST .....	125	ONUREG .....	143
MEKTOVI .....	25	OPSUMIT .....	161
<i>mifepristone oral tablet 300 mg .....</i>	126	ORENITRAM .....	161
<i>miglustat .....</i>	127	ORENITRAM MONTH 1 .....	161
MODEYSO .....	128	ORENITRAM MONTH 2 .....	161
MOUNJARO SUBCUTANEOUS SOLUTION		ORENITRAM MONTH 3 .....	161
AUTO-INJECTOR 10 MG/0.5ML, 12.5		ORIAHNN .....	144
MG/0.5ML, 15 MG/0.5ML, 2.5 MG/0.5ML,		ORKAMBI .....	36
5 MG/0.5ML, 7.5 MG/0.5ML .....	75	<i>ormalvi .....</i>	51
MYTESI .....	129	ORSERDU .....	145
NAYZILAM .....	193	OTEZLA ORAL TABLET 30 MG .....	21
NERLYNX .....	130	OTEZLA ORAL TABLET THERAPY PACK 10 &	
NEXLETOL .....	15	20 & 30 MG .....	21
NEXLIZET .....	15	OZEMPIC (0.25 OR 0.5 MG/DOSE)	
<i>nilotinib hcl .....</i>	131	SUBCUTANEOUS SOLUTION PEN-INJECTOR	
NINLARO .....	132	2 MG/3ML .....	75
<i>nitisinone .....</i>	133	OZEMPIC (1 MG/DOSE) SUBCUTANEOUS	
NORDITROPIN FLEXPOR SUBCUTANEOUS		SOLUTION PEN-INJECTOR 4 MG/3ML .....	75
SOLUTION PEN-INJECTOR .....	78	OZEMPIC (2 MG/DOSE) .....	75
NUBEQA .....	160	PANRETIN .....	147
NUCALA SUBCUTANEOUS SOLUTION		<i>pazopanib hcl .....</i>	148
AUTO-INJECTOR .....	134	PEGASYS SUBCUTANEOUS SOLUTION 180	
NUCALA SUBCUTANEOUS SOLUTION		MCG/ML .....	193
PREFILLED SYRINGE 100 MG/ML .....	134	PEGASYS SUBCUTANEOUS SOLUTION	
NUCALA SUBCUTANEOUS SOLUTION		PREFILLED SYRINGE .....	193
RECONSTITUTED .....	134	PEMAZYRE ORAL TABLET 13.5 MG, 4.5	
NUDEXTA .....	136	MG, 9 MG .....	150
NUPLAZID ORAL CAPSULE .....	137	PIQRAY (200 MG DAILY DOSE) .....	151
NUPLAZID ORAL TABLET 10 MG .....	137	PIQRAY (250 MG DAILY DOSE) .....	151
NURTEC .....	37	PIQRAY (300 MG DAILY DOSE) .....	151
		<i>pirfenidone oral tablet 267 mg, 801 mg ....</i>	152
		POMALYST .....	153
		PRETOMANID .....	154

PREVYMIS ORAL TABLET .....	155	SECUADO TRANSDERMAL PATCH 24 HOUR	
PRIVIGEN .....	88	3.8 MG/24HR, 5.7 MG/24HR, 7.6	
PROLASTIN-C INTRAVENOUS SOLUTION ....	157	MG/24HR .....	181
<i>promethazine hcl oral solution 6.25</i>		SIGNIFOR .....	184
<i>mg/5ml .....</i>	159	<i>sildenafil citrate oral suspension</i>	
<i>promethazine hcl oral tablet .....</i>	159	<i>reconstituted .....</i>	161
<i>promethazine hcl rectal suppository 12.5</i>		<i>sildenafil citrate oral tablet 20 mg .....</i>	161
<i>mg, 25 mg .....</i>	159	<i>simvastatin oral tablet 80 mg .....</i>	185
PROMETHEGAN RECTAL SUPPOSITORY 25		SIRTURO .....	186
MG, 50 MG .....	159	SIVEXTRO .....	187
QINLOCK .....	163	SKYCLARYS .....	188
QULIPTA .....	37	<i>sodium oxybate .....</i>	146
RECORLEV .....	164	<i>sofosbuvir-velpatasvir .....</i>	189
REPATHA .....	166	SOMAVERT .....	190
REPATHA PUSHTRONEX SYSTEM .....	166	<i>sorafenib tosylate .....</i>	192
REPATHA SURECLICK .....	166	SOTYKTU .....	21
<i>retacrit injection solution 10000 unit/ml,</i>		STELARA SUBCUTANEOUS SOLUTION 45	
<i>2000 unit/ml, 20000 unit/ml, 3000 unit/ml,</i>		MG/0.5ML .....	21
<i>4000 unit/ml, 40000 unit/ml .....</i>	168	STELARA SUBCUTANEOUS SOLUTION	
RETEVMO ORAL TABLET 120 MG, 160 MG,		PREFILLED SYRINGE .....	21
40 MG, 80 MG .....	169	STIVARGA .....	194
REVCovi .....	170	SUCRAID .....	195
REVUFORJ ORAL TABLET 110 MG, 160 MG,		<i>sunitinib malate .....</i>	196
25 MG .....	171	SUNOSI ORAL TABLET 150 MG, 75 MG .....	197
REXULTI ORAL TABLET 0.25 MG, 0.5 MG, 1		SYMDEKO .....	36
MG, 2 MG, 3 MG, 4 MG .....	181	TABRECTA .....	198
REZDIFFRA .....	172	<i>tadalafil (pah) .....</i>	161
REZLIDHIA .....	173	<i>tadalafil oral tablet 2.5 mg, 5 mg .....</i>	199
REZUROCK .....	174	TAFINLAR .....	125
ROMVIMZA .....	175	TAGRISSO .....	202
ROZLYTREK .....	176	TALZENNA ORAL CAPSULE 0.1 MG, 0.25	
RUBRACA .....	177	MG, 0.35 MG, 0.5 MG, 0.75 MG, 1 MG .....	203
RYBELSUS (FORMULATION R2) ORAL		<i>tasimelteon .....</i>	204
TABLET 1.5 MG, 4 MG, 9 MG .....	75	<i>tazarotene external cream .....</i>	214
RYBELSUS ORAL TABLET 14 MG, 3 MG, 7		TAZVERIK .....	205
MG .....	75	TEPMETKO .....	206
RYDAPT .....	178	<i>teriparatide subcutaneous solution pen-</i>	
<i>sajazir subcutaneous solution prefilled</i>		<i>injector 560 mcg/2.24ml .....</i>	207
<i>syringe .....</i>	79	<i>testosterone transdermal gel 20.25 mg/act</i>	
<i>sapropterin dihydrochloride oral packet ....</i>	179	<i>(1.62%) .....</i>	209
<i>sapropterin dihydrochloride oral tablet ....</i>	179	TIBSOVO .....	210
SCEMBLIX ORAL TABLET 100 MG, 20 MG,		<i>tolvaptan oral tablet 15 mg, 30 mg .....</i>	211
40 MG .....	180		

<i>tolvaptan oral tablet 15 mg</i> <i>tolvaptan (hyponatremia), 30 mg</i> <i>tolvaptan (hyponatremia)</i> .....	213	VITRAKVI ORAL CAPSULE 100 MG, 25 MG .	228
<i>tolvaptan oral tablet therapy pack 15 mg, 30 &amp; 15 mg, 45 &amp; 15 mg, 60 &amp; 30 mg, 90 &amp; 30 mg</i> .....	211	VITRAKVI ORAL SOLUTION .....	228
<i>torpenz</i> .....	64	VIVJOA .....	229
<i>tretinoin external cream</i> .....	214	VIZIMPRO ORAL TABLET 15 MG, 30 MG, 45 MG .....	230
<i>tretinoin external gel 0.01 %, 0.025 %</i> .....	214	VONJO .....	231
<i>tridacaine ii</i> .....	113	VORANIGO ORAL TABLET 10 MG, 40 MG ...	232
TRIKAFTA ORAL TABLET THERAPY PACK 100-50-75 & 150 MG, 50-25-37.5 & 75 MG .	36	VOWST .....	233
TRIKAFTA ORAL THERAPY PACK.....	36	VYNDAMAX .....	200
TRULICITY SUBCUTANEOUS SOLUTION AUTO-INJECTOR.....	75	VYNDAQEL.....	200
TRUQAP ORAL TABLET .....	215	WELIREG .....	234
TRUQAP ORAL TABLET THERAPY PACK .....	215	WINREVAIR SUBCUTANEOUS KIT 2 X 45 MG, 2 X 60 MG, 45 MG, 60 MG .....	235
TUKYSA ORAL TABLET 150 MG, 50 MG .....	216	XALKORI .....	237
TURALIO ORAL CAPSULE 125 MG .....	217	XDEMVI .....	238
<i>tyenne subcutaneous</i> .....	21	XERMELO .....	239
UBRELVY .....	37	XIFAXAN ORAL TABLET 550 MG .....	240
UDENYCA .....	149	XOLAIR.....	242
UDENYCA ONBODY .....	149	XOSPATA.....	244
<i>ustekinumab subcutaneous</i> .....	21	XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 50 MG .....	245
VALCHLOR .....	218	XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 10 MG, 40 MG .....	245
VALTOCO 10 MG DOSE .....	193	XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 40 MG .....	245
VALTOCO 15 MG DOSE NASAL LIQUID THERAPY PACK 2 X 7.5 MG/0.1ML.....	193	XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG .....	245
VALTOCO 20 MG DOSE NASAL LIQUID THERAPY PACK 2 X 10 MG/0.1ML.....	193	XPOVIO (60 MG TWICE WEEKLY) .....	245
VALTOCO 5 MG DOSE .....	193	XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG .....	245
VANFLYTA .....	219	XPOVIO (80 MG TWICE WEEKLY) .....	245
VANRAFIA .....	220	XTANDI .....	160
VELSIPITY .....	21	XURIDEN .....	246
VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG .....	222	<i>yargesa</i> .....	127
VENCLEXTA STARTING PACK .....	222	<i>zaleplon</i> .....	183
VERQUVO ORAL TABLET 10 MG, 2.5 MG, 5 MG .....	223	ZEJULA ORAL TABLET .....	247
VERZENIO .....	225	ZELBORAF .....	43
VIBERZI .....	226	<i>zolpidem tartrate oral tablet</i> .....	183
VIJOICE ORAL PACKET .....	227	ZTALMY .....	248
VIJOICE ORAL TABLET THERAPY PACK 125 MG, 200 & 50 MG, 50 MG .....	227	ZURZUVAE .....	249
		ZYDELIG .....	250
		ZYKADIA ORAL TABLET .....	251