

## QUICK GUIDE: MAT Use for Opioid Use Disorder



Implant



Pill










Patch



Liquid



Injection

	<b>BUPRENORPHINE</b>	<b>METHADONE</b>	<b>NALTREXONE</b>
<b>COMMON BRANDS</b>	Suboxone, Zubsolv, Bunavail, Subutex, Probuphine, Sublocade	Methadose, Diskets, Dolophine	Vivitrol
<b>TYPE</b>	   	 	
<b>HOW IT WORKS</b>	<ul style="list-style-type: none"> <li>• Makes the brain think it is still getting the problem opioid. Prevents cravings and withdrawal symptoms and reduces the risk of overdose.</li> <li>• Buprenorphine can be prescribed by a trained provider in a doctor's office or other health care setting, as well as in a narcotic treatment program (NTP).</li> </ul>	<ul style="list-style-type: none"> <li>• Makes the brain think it is still getting the problem opioid. Prevents cravings and withdrawal symptoms and reduces the risk of overdose.</li> <li>• Methadone is dispensed only in highly regulated NTPs.</li> </ul>	<ul style="list-style-type: none"> <li>• Blocks the effects of opioids.</li> <li>• Naltrexone is not a controlled substance and can be prescribed or administered in any health care or substance use disorder (SUD) setting, such as a doctor's office or clinic.</li> </ul>
<b>THINGS TO CONSIDER</b>	<ul style="list-style-type: none"> <li>• Treatment can start quickly, as soon as someone enters withdrawal.</li> <li>• Flexible dosing schedule.</li> <li>• Relapse risk increases if you forget or choose not to take medication.</li> <li>• Common side effects are headache, nausea, and constipation.</li> </ul>	<ul style="list-style-type: none"> <li>• Treatment can start right away, no need for detoxification.</li> <li>• Less flexible schedule. Dosing occurs in the early morning.</li> <li>• Side effects include constipation, sexual problems, swelling, and sweating and potential heart problems.</li> </ul>	<ul style="list-style-type: none"> <li>• Less evidence for effectiveness in OUD treatment than buprenorphine or methadone.</li> <li>• Does not cause physical dependence.</li> <li>• Not recommended for pregnant women as detox can harm the baby. Methadone or buprenorphine are recommended for pregnant women with OUD.</li> </ul>

CONTINUED ON BACK

## QUICK GUIDE: MAT Use for Opioid Use Disorder

CONTINUED FROM FRONT	BUPRENORPHINE	METHADONE	NALTREXONE
<b>THINGS TO CONSIDER</b>	<ul style="list-style-type: none"> <li>• Causes physical dependence. If or when you want to come off the drug, you will need to do so slowly to minimize the discomfort of detox symptoms.</li> <li>• Buprenorphine is sometimes used short term to relieve pain associated with detox, but more often used long term, known as maintenance treatment.</li> </ul>	<ul style="list-style-type: none"> <li>• Causes physical dependence. If you want to come off the drug, you will need to do so slowly to minimize the discomfort of detox symptoms.</li> <li>• May cause drowsiness at first before maintenance dose is determined.</li> <li>• Methadone is often a good option for people who have used opioids for a long time or have been unsuccessful with other treatments.</li> </ul>	<ul style="list-style-type: none"> <li>• Detox from opioids is required before taking naltrexone. Side effects may include stomach pain, nausea, vomiting, headache, joint pain, trouble sleeping and anxiety. Some people also report soreness in the area of the injection.</li> <li>• Injection form of the medication lasts for about 30 days before it wears off. Overdose risk can be higher after naltrexone wears off due to decrease in tolerance.</li> </ul>
<b>QUESTIONS</b>	<ul style="list-style-type: none"> <li>• Can you commit to taking this medication daily?</li> <li>• Are you comfortable with taking a medication that requires time to taper off to minimize the discomfort of detox?</li> </ul>	<ul style="list-style-type: none"> <li>• Have you used opioids for a long time, or have you found other treatments have not worked well for you?</li> <li>• Can you come to the clinic in the early morning for dosing? Will you need to make arrangements for work or transportation?</li> <li>• Do you work in an industry with heavy machinery? Could your work be affected by possible drowsiness during your initial dosing period?</li> <li>• Are you comfortable with taking a medication that requires time to taper off to minimize the discomfort of detox?</li> </ul>	<ul style="list-style-type: none"> <li>• Have you detoxed from opioids, or would you be willing to detox to take this medication?</li> <li>• Can you commit to making an appointment once every month to continue receiving the injection?</li> <li>• Do you have any medical needs that would be affected by blocking the opioid receptors? For example, do you use opioids to reduce chronic pain?</li> </ul>